

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number  
WO 01/79472 A2

- (51) International Patent Classification<sup>7</sup>: C12N 9/00 (74) Agent: MARSH, David, R.; Arnold & Porter, 555 12th Street, NW, Washington, DC 20004-1206 (US).
- (21) International Application Number: PCT/US01/12334
- (22) International Filing Date: 13 April 2001 (13.04.2001) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/549,848 14 April 2000 (14.04.2000) US  
09/688,069 14 October 2000 (14.10.2000) US
- (71) Applicant: MONSANTO TECHNOLOGY LLP  
[US/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US).
- (72) Inventors: SUBRAMANIAM, Sai, S.; 33 Dinsmore Avenue #304, Framingham, MA 01702 (US). SLATER, Steven, C.; 21 Brucewood Road, Acton, MA 01720 (US). KARBERG, Katherine; 37 Regent Street #3, Cambridge, MA 02140 (US). CHEN, Ridong; 11131-I Westport Station Drive, Maryland Heights, MO 63043 (US). VALENTIN, Henry, E.; 37 Regent Street #3, Cambridge, MA 02140 (US). WONG, Yun-Hua, Huang; 14043 Forest Crest Drive, Chesterfield, MO 63017 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/79472 A2

(54) Title: NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN TOCOPHEROL SYNTHESIS

(57) Abstract: Nucleic acid sequences and methods are provided for producing plants and seeds having altered tocopherol content and compositions. The methods find particular use in increasing the tocopherol levels in plants, and in providing desirable tocopherol compositions in a host plant cell.

5                   **NUCLEIC ACID SEQUENCES TO PROTEINS INVOLVED IN  
                          TOCOPHEROL SYNTHESIS**

**INTRODUCTION**

**TECHNICAL FIELD**

10           The present invention is directed to nucleic acid and amino acid sequences and constructs, and methods related thereto.

**BACKGROUND**

          Isoprenoids are ubiquitous compounds found in all living organisms. Plants synthesize a diverse array of greater than 22,000 isoprenoids (Connolly and Hill  
15   (1992) *Dictionary of Terpenoids*, Chapman and Hall, New York, NY). In plants, isoprenoids play essential roles in particular cell functions such as production of sterols, contributing to eukaryotic membrane architecture, acyclic polyprenoids found in the side chain of ubiquinone and plastoquinone, growth regulators like abscisic acid, gibberellins, brassinosteroids or the photosynthetic pigments chlorophylls and  
20   carotenoids. Although the physiological role of other plant isoprenoids is less evident, like that of the vast array of secondary metabolites, some are known to play key roles mediating the adaptative responses to different environmental challenges. In spite of the remarkable diversity of structure and function, all isoprenoids originate from a single metabolic precursor, isopentenyl diphosphate (IPP) (Wright, (1961) *Annu. Rev. Biochem.* 20:525-548; and Spurgeon and Porter, (1981) in Biosynthesis of Isoprenoid  
25   Compounds., Porter and Spurgeon eds (John Wiley, New York) Vol. 1, pp1-46).

          A number of unique and interconnected biochemical pathways derived from the isoprenoid pathway leading to secondary metabolites, including tocopherols, exist in chloroplasts of higher plants. Tocopherols not only perform vital functions in

plants, but are also important from mammalian nutritional perspectives. In plastids, tocopherols account for up to 40% of the total quinone pool.

Tocopherols and tocotrienols (unsaturated tocopherol derivatives) are well known antioxidants, and play an important role in protecting cells from free radical damage, and in the prevention of many diseases, including cardiac disease, cancer, cataracts, retinopathy, Alzheimer's disease, and neurodegeneration, and have been shown to have beneficial effects on symptoms of arthritis, and in anti-aging. Vitamin E is used in chicken feed for improving the shelf life, appearance, flavor, and oxidative stability of meat, and to transfer tocopherols from feed to eggs. Vitamin E has been shown to be essential for normal reproduction, improves overall performance, and enhances immunocompetence in livestock animals. Vitamin E supplement in animal feed also imparts oxidative stability to milk products.

The demand for natural tocopherols as supplements has been steadily growing at a rate of 10-20% for the past three years. At present, the demand exceeds the supply for natural tocopherols, which are known to be more biopotent than racemic mixtures of synthetically produced tocopherols. Naturally occurring tocopherols are all *d*-stereoisomers, whereas synthetic  $\alpha$ -tocopherol is a mixture of eight *d,l*- $\alpha$ -tocopherol isomers, only one of which (12.5%) is identical to the natural *d*- $\alpha$ -tocopherol. Natural *d*- $\alpha$ -tocopherol has the highest vitamin E activity (1.49 IU/mg) when compared to other natural tocopherols or tocotrienols. The synthetic  $\alpha$ -tocopherol has a vitamin E activity of 1.1 IU/mg. In 1995, the worldwide market for raw refined tocopherols was \$1020 million; synthetic materials comprised 85-88% of the market, the remaining 12-15% being natural materials. The best sources of natural tocopherols and tocotrienols are vegetable oils and grain products. Currently, most of the natural Vitamin E is produced from  $\gamma$ -tocopherol derived from soy oil processing, which is subsequently converted to  $\alpha$ -tocopherol by chemical modification ( $\alpha$ -tocopherol exhibits the greatest biological activity).

Methods of enhancing the levels of tocopherols and tocotrienols in plants, especially levels of the more desirable compounds that can be used directly, without

chemical modification, would be useful to the art as such molecules exhibit better functionality and bioavailability.

In addition, methods for the increased production of other isoprenoid derived compounds in a host plant cell is desirable. Furthermore, methods for the production of particular isoprenoid compounds in a host plant cell is also needed.

#### SUMMARY OF THE INVENTION

The present invention is directed to sequences to proteins involved in tocopherol synthesis. The polynucleotides and polypeptides of the present invention include those derived from prokaryotic and eukaryotic sources.

Thus, one aspect of the present invention relates to prenyltransferase, and in particular to isolated polynucleotide sequences encoding prenyltransferase proteins and polypeptides related thereto. In particular, isolated nucleic acid sequences encoding prenyltransferase proteins from bacterial and plant sources are provided.

In another aspect, the present invention provides isolated polynucleotide sequences encoding tocopherol cyclase, and polypeptides related thereto. In particular, isolated nucleic acid sequences encoding tocopherol cyclase proteins from bacterial and plant sources are provided.

Another aspect of the present invention relates to oligonucleotides which include partial or complete prenyltransferase or tocopherol cyclase encoding sequences.

It is also an aspect of the present invention to provide recombinant DNA constructs which can be used for transcription or transcription and translation (expression) of prenyltransferase or tocopherol cyclase. In particular, constructs are provided which are capable of transcription or transcription and translation in host cells.

In another aspect of the present invention, methods are provided for production of prenyltransferase or tocopherol cyclase in a host cell or progeny thereof. In particular, host cells are transformed or transfected with a DNA construct which can be used for transcription or transcription and translation of

prenyltransferase or tocopherol cyclase. The recombinant cells which contain prenyltransferase or tocopherol cyclase are also part of the present invention.

In a further aspect, the present invention relates to methods of using polynucleotide and polypeptide sequences to modify the tocopherol content of host  
5 cells, particularly in host plant cells. Plant cells having such a modified tocopherol content are also contemplated herein. Methods and cells in which both prenyltransferase and tocopherol cyclase are expressed in a host cell are also part of the present invention.

The modified plants, seeds and oils obtained by the expression of the  
10 prenyltransferase or tocopherol cyclase are also considered part of the invention.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 provides an amino acid sequence alignment between ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 are performed using ClustalW.

Figure 2 provides a schematic picture of the expression construct pCGN10800.  
15 Figure 3 provides a schematic picture of the expression construct pCGN10801.  
Figure 4 provides a schematic picture of the expression construct pCGN10803.  
Figure 5 provides a schematic picture of the construct pCGN10806.  
Figure 6 provides a schematic picture of the construct pCGN10807.  
Figure 7 provides a schematic picture of the construct pCGN10808.  
20 Figure 8 provides a schematic picture of the expression construct pCGN10809.  
Figure 9 provides a schematic picture of the expression construct pCGN10810.  
Figure 10 provides a schematic picture of the expression construct pCGN10811.  
Figure 11 provides a schematic picture of the expression construct pCGN10812.  
Figure 12 provides a schematic picture of the expression construct pCGN10813.  
25 Figure 13 provides a schematic picture of the expression construct pCGN10814.  
Figure 14 provides a schematic picture of the expression construct pCGN10815.  
Figure 15 provides a schematic picture of the expression construct pCGN10816.  
Figure 16 provides a schematic picture of the expression construct pCGN10817.  
Figure 17 provides a schematic picture of the expression construct pCGN10819.

Figure 18 provides a schematic picture of the expression construct pCGN10824.

Figure 19 provides a schematic picture of the expression construct pCGN10825.

Figure 20 provides a schematic picture of the expression construct pCGN10826.

Figure 21 provides an amino acid sequence alignment using ClustalW between the  
5 *Synechocystis* prenyltransferase sequences.

Figure 22 provides an amino acid sequence of the ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 protein sequences from *Arabidopsis* and the slr1736, slr0926, slr11899, slr0056, and the slr1518 amino acid sequences from *Synechocystis*.

Figure 23 provides the results of the enzymatic assay from preparations of  
10 wild type *Synechocystis* strain 6803, and *Synechocystis* slr1736 knockout.

Figure 24 provides bar graphs of HPLC data obtained from seed extracts of transgenic *Arabidopsis* containing pCGN10822, which provides of the expression of the ATPT2 sequence, in the sense orientation, from the napin promoter. Provided are graphs for alpha, gamma, and delta tocopherols, as well as total tocopherol for 22  
15 transformed lines, as well as a nontransformed (wildtype) control.

Figure 25 provides a bar graph of HPLC analysis of seed extracts from *Arabidopsis* plants transformed with pCGN10803 (35S-ATPT2, in the antisense orientation), pCGN10822 (line 1625, napin ATPT2 in the sense orientation), pCGN10809 (line 1627, 35S-ATPT3 in the sense orientation), a nontransformed (wt)  
20 control, and an empty vector transformed control.

Figure 26 shows total tocopherol levels measured in T# *Arabidopsis* seed of line.

Figure 27 shows total tocopherol levels measured in T# *Arabidopsis* seed of line.

Figure 28 shows total tocopherol levels measured in developing canola seed of  
25 line 10822-1.

Figure 29: shows results of phytol prenyltransferase activity assay using *Synechocystis* wild type and slr1737 knockout mutant membrane preparations.



Figure 30 is the chromatograph from an HPLC analysis of *Synechocystis* extracts.

Figure 31 is a sequence alignment of the *Arabidopsis* homologue with the sequence of the public database.

5        Figure 32 shows the results of hydropathic analysis of slr1737

Figure 33 shows the results of hydropathic analysis of the *Arabidopsis* homologue of slr1737.

Figure 34 shows the catalytic mechanism of various cyclase enzymes

10        Figure 35 is a sequence alignment of slr1737, slr1737 *Arabidopsis* homologue and the *Arabidopsis* chalcone isomerase.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides, *inter alia*, compositions and methods for altering (for example, increasing and decreasing) the tocopherol levels and/or modulating their ratios in host cells. In particular, the present invention provides  
15        polynucleotides, polypeptides, and methods of use thereof for the modulation of tocopherol content in host plant cells.

The biosynthesis of  $\alpha$ -tocopherol in higher plants involves condensation of homogentisic acid and phytylpyrophosphate to form 2-methyl-6 phytylbenzoquinol that can, by cyclization and subsequent methylations (Fiedler et al., 1982, *Planta*, 155:  
20        511-515, Soll et al., 1980, *Arch. Biochem. Biophys.* 204: 544-550, Marshall et al., 1985 *Phytochem.*, 24: 1705-1711, all of which are herein incorporated by reference in their entirety), form various tocopherols.

The *Arabidopsis pds2* mutant identified and characterized by Norris *et al.* (1995), is deficient in tocopherol and plastoquinone-9 accumulation. Further genetic  
25        and biochemical analysis suggested that the protein encoded by *PDS2* may be responsible for the prenylation of homogentisic acid. The *PDS2* locus identified by Norris *et al.* (1995) has been hypothesized to possibly encode the tocopherol phytyl-prenyltransferase, as the *pds2* mutant fails to accumulate tocopherols.

Norris *et al.* (1995) determined that in *Arabidopsis pds2* lies at the top of chromosome 3, approximately 7 centimorgans above long hypocotyl2, based on the genetic map. ATPT2 is located on chromosome 2 between 36 and 41 centimorgans, lying on BAC F19F24, indicating that ATPT2 does not correspond to *PDS2*. Thus, it is an aspect of the present invention to provide novel polynucleotides and polypeptides involved in the prenylation of homogentisic acid. This reaction may be a rate limiting step in tocopherol biosynthesis, and this gene has yet to be isolated.

U.S. Patent No. 5,432,069 describes the partial purification and characterization of tocopherol cyclase from *Chlorella protothecoides*, *Dunaliella salina* and wheat. The cyclase described as being glycine rich, water soluble and with a predicted MW of 48-50kDa. However, only limited peptide fragment sequences were available.

In one aspect, the present invention provides polynucleotide and polypeptide sequences involved in the prenylation of straight chain and aromatic compounds. Straight chain prenyltransferases as used herein comprises sequences which encode proteins involved in the prenylation of straight chain compounds, including, but not limited to, geranyl geranyl pyrophosphate and farnesyl pyrophosphate. Aromatic prenyltransferases, as used herein, comprises sequences which encode proteins involved in the prenylation of aromatic compounds, including, but not limited to, menaquinone, ubiquinone, chlorophyll, and homogentisic acid. The prenyltransferase of the present invention preferably prenylates homogentisic acid.

In another aspect, the invention provides polynucleotide and polypeptide sequences to tocopherol cyclization enzymes. The 2,3-dimethyl-5-phytylplastoquinol cyclase (tocopherol cyclase) is responsible for the cyclization of 2,3-dimethyl-5-phytylplastoquinol to tocopherol.

#### **Isolated Polynucleotides, Proteins, and Polypeptides**

A first aspect of the present invention relates to isolated prenyltransferase polynucleotides. Another aspect of the present invention relates to isolated tocopherol cyclase polynucleotides. The polynucleotide sequences of the present invention

include isolated polynucleotides that encode the polypeptides of the invention having a deduced amino acid sequence selected from the group of sequences set forth in the Sequence Listing and to other polynucleotide sequences closely related to such sequences and variants thereof.

5           The invention provides a polynucleotide sequence identical over its entire length to each coding sequence as set forth in the Sequence Listing. The invention also provides the coding sequence for the mature polypeptide or a fragment thereof, as well as the coding sequence for the mature polypeptide or a fragment thereof in a reading frame with other coding sequences, such as those encoding a leader or  
10   secretory sequence, a pre-, pro-, or prepro- protein sequence. The polynucleotide can also include non-coding sequences, including for example, but not limited to, non-coding 5' and 3' sequences, such as the transcribed, untranslated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence that encodes additional  
15   amino acids. For example, a marker sequence can be included to facilitate the purification of the fused polypeptide. Polynucleotides of the present invention also include polynucleotides comprising a structural gene and the naturally associated sequences that control gene expression.

The invention also includes polynucleotides of the formula:

20                            $X-(R_1)_n-(R_2)-(R_3)_n-Y$

wherein, at the 5' end, X is hydrogen, and at the 3' end, Y is hydrogen or a metal,  $R_1$  and  $R_3$  are any nucleic acid residue, n is an integer between 1 and 3000, preferably between 1 and 1000 and  $R_2$  is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from the group set forth in the Sequence Listing and  
25   preferably those of SEQ ID NOs: 1, 3, 5, 7, 8, 10, 11, 13-16, 18, 23, 29, 36, and 38. In the formula,  $R_2$  is oriented so that its 5' end residue is at the left, bound to  $R_1$ , and its 3' end residue is at the right, bound to  $R_3$ . Any stretch of nucleic acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

The invention also relates to variants of the polynucleotides described herein that encode for variants of the polypeptides of the invention. Variants that are fragments of the polynucleotides of the invention can be used to synthesize full-length polynucleotides of the invention. Preferred embodiments are polynucleotides  
5 encoding polypeptide variants wherein 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues of a polypeptide sequence of the invention are substituted, added or deleted, in any combination. Particularly preferred are substitutions, additions, and deletions that are silent such that they do not alter the properties or activities of the polynucleotide or polypeptide.

10 Further preferred embodiments of the invention that are at least 50%, 60%, or 70% identical over their entire length to a polynucleotide encoding a polypeptide of the invention, and polynucleotides that are complementary to such polynucleotides. More preferable are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding a polypeptide of the  
15 invention and polynucleotides that are complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length are particularly preferred, those at least 95% identical are especially preferred. Further, those with at least 97% identity are highly preferred and those with at least 98% and 99% identity are particularly highly preferred, with those at least 99% being the most highly  
20 preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptides encoded by the polynucleotides set forth in the Sequence Listing.

The invention further relates to polynucleotides that hybridize to the above-  
25 described sequences. In particular, the invention relates to polynucleotides that hybridize under stringent conditions to the above-described polynucleotides. As used herein, the terms "stringent conditions" and "stringent hybridization conditions" mean that hybridization will generally occur if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization

conditions is overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/milliliter denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at approximately 65°C. Other hybridization and wash conditions are well known and are exemplified in Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, NY (1989), particularly Chapter 11.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in the Sequence Listing under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers as described herein.

As discussed herein regarding polynucleotide assays of the invention, for example, polynucleotides of the invention can be used as a hybridization probe for RNA, cDNA, or genomic DNA to isolate full length cDNAs or genomic clones encoding a polypeptide and to isolate cDNA or genomic clones of other genes that have a high sequence similarity to a polynucleotide set forth in the Sequence Listing. Such probes will generally comprise at least 15 bases. Preferably such probes will have at least 30 bases and can have at least 50 bases. Particularly preferred probes will have between 30 bases and 50 bases, inclusive.

The coding region of each gene that comprises or is comprised by a polynucleotide sequence set forth in the Sequence Listing may be isolated by screening using a DNA sequence provided in the Sequence Listing to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to identify members of the library which hybridize to the probe. For

example, synthetic oligonucleotides are prepared which correspond to the prenyltransferase or tocopherol cyclase EST sequences. The oligonucleotides are used as primers in polymerase chain reaction (PCR) techniques to obtain 5' and 3' terminal sequence of prenyltransferase or tocopherol cyclase genes. Alternatively, 5 where oligonucleotides of low degeneracy can be prepared from particular prenyltransferase or tocopherol cyclase peptides, such probes may be used directly to screen gene libraries for prenyltransferase or tocopherol cyclase gene sequences. In particular, screening of cDNA libraries in phage vectors is useful in such methods due to lower levels of background hybridization.

10 Typically, a prenyltransferase or tocopherol cyclase sequence obtainable from the use of nucleic acid probes will show 60-70% sequence identity between the target prenyltransferase or tocopherol cyclase sequence and the encoding sequence used as a probe. However, lengthy sequences with as little as 50-60% sequence identity may also be obtained. The nucleic acid probes may be a lengthy fragment of the nucleic 15 acid sequence, or may also be a shorter, oligonucleotide probe. When longer nucleic acid fragments are employed as probes (greater than about 100 bp), one may screen at lower stringencies in order to obtain sequences from the target sample which have 20-50% deviation (i.e., 50-80% sequence homology) from the sequences used as probe. Oligonucleotide probes can be considerably shorter than the entire nucleic acid 20 sequence encoding an prenyltransferase or tocopherol cyclase enzyme, but should be at least about 10, preferably at least about 15, and more preferably at least about 20 nucleotides. A higher degree of sequence identity is desired when shorter regions are used as opposed to longer regions. It may thus be desirable to identify regions of highly conserved amino acid sequence to design oligonucleotide probes for detecting 25 and recovering other related prenyltransferase or tocopherol cyclase genes. Shorter probes are often particularly useful for polymerase chain reactions (PCR), especially when highly conserved sequences can be identified. (See, Gould, *et al.*, *PNAS USA* (1989) 86:1934-1938.).

Another aspect of the present invention relates to prenyltransferase or tocopherol cyclase polypeptides. Such polypeptides include isolated polypeptides set forth in the Sequence Listing, as well as polypeptides and fragments thereof, particularly those polypeptides which exhibit prenyltransferase or tocopherol cyclase activity and also those polypeptides which have at least 50%, 60% or 70% identity, preferably at least 80% identity, more preferably at least 90% identity, and most preferably at least 95% identity to a polypeptide sequence selected from the group of sequences set forth in the Sequence Listing, and also include portions of such polypeptides, wherein such portion of the polypeptide preferably includes at least 30 amino acids and more preferably includes at least 50 amino acids.

"Identity", as is well understood in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods including, but not limited to, those described in *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M. and Griffin, H.G., eds., Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press (1987); *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., Stockton Press, New York (1991); and Carillo, H., and Lipman, D., *SIAM J Applied Math*, 48:1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available programs. Computer programs which can be used to determine identity between two sequences include, but are not limited to, GCG (Devereux, J., et al., *Nucleic Acids Research* 12(1):387 (1984); suite of five BLAST programs, three designed for nucleotide sequences queries (BLASTN, BLASTX, and TBLASTX) and two

designed for protein sequence queries (BLASTP and TBLASTN) (Coulson, *Trends in Biotechnology*, 12: 76-80 (1994); Birren, *et al.*, *Genome Analysis*, 1: 543-559 (1997)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., *et al.*, NCBI NLM NIH, Bethesda, MD 20894; Altschul, S., *et al.*, *J. Mol. Biol.*, 215:403-410 (1990)). The well known Smith Waterman algorithm can also be used to determine identity.

Parameters for polypeptide sequence comparison typically include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)  
10 Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci USA* 89:10915-10919 (1992)

Gap Penalty: 12

Gap Length Penalty: 4

A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above  
15 parameters along with no penalty for end gap are the default parameters for peptide comparisons.

Parameters for polynucleotide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)

20 Comparison matrix: matches = +10; mismatches = 0

Gap Penalty: 50

Gap Length Penalty: 3

A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above  
25 parameters are the default parameters for nucleic acid comparisons.

The invention also includes polypeptides of the formula:



wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal,  $R_1$  and  $R_3$  are any amino acid residue,  $n$  is an integer between 1



and 1000, and  $R_2$  is an amino acid sequence of the invention, particularly an amino acid sequence selected from the group set forth in the Sequence Listing and preferably those encoded by the sequences provided in SEQ ID NOs: 2, 4, 6, 9, 12, 17, 19-22, 24-28, 30, 32-35, 37, and 39. In the formula,  $R_2$  is oriented so that its amino terminal residue is at the left, bound to  $R_1$ , and its carboxy terminal residue is at the right, bound to  $R_3$ . Any stretch of amino acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

Polypeptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising a sequence selected from the group of a sequence contained in the Sequence Listing set forth herein .

The polypeptides of the present invention can be mature protein or can be part of a fusion protein.

Fragments and variants of the polypeptides are also considered to be a part of the invention. A fragment is a variant polypeptide which has an amino acid sequence that is entirely the same as part but not all of the amino acid sequence of the previously described polypeptides. The fragments can be "free-standing" or comprised within a larger polypeptide of which the fragment forms a part or a region, most preferably as a single continuous region. Preferred fragments are biologically active fragments which are those fragments that mediate activities of the polypeptides of the invention, including those with similar activity or improved activity or with a decreased activity. Also included are those fragments that antigenic or immunogenic in an animal, particularly a human.

Variants of the polypeptide also include polypeptides that vary from the sequences set forth in the Sequence Listing by conservative amino acid substitutions, substitution of a residue by another with like characteristics. In general, such substitutions are among Ala, Val, Leu and Ile; between Ser and Thr; between Asp and Glu; between Asn and Gln; between Lys and Arg; or between Phe and Tyr.

Particularly preferred are variants in which 5 to 10; 1 to 5; 1 to 3 or one amino acid(s) are substituted, deleted, or added, in any combination.

5 Variants that are fragments of the polypeptides of the invention can be used to produce the corresponding full length polypeptide by peptide synthesis. Therefore, these variants can be used as intermediates for producing the full-length polypeptides of the invention.

The polynucleotides and polypeptides of the invention can be used, for example, in the transformation of host cells, such as plant host cells, as further discussed herein.

10 The invention also provides polynucleotides that encode a polypeptide that is a mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids within the mature polypeptide (for example, when the mature form of the protein has more than one polypeptide chain). Such sequences can, for example, play a role in the processing of a protein from a precursor to a mature form, allow protein  
15 transport, shorten or lengthen protein half-life, or facilitate manipulation of the protein in assays or production. It is contemplated that cellular enzymes can be used to remove any additional amino acids from the mature protein.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. The inactive  
20 precursors generally are activated when the prosequences are removed. Some or all of the prosequences may be removed prior to activation. Such precursor protein are generally called proproteins.

#### **Plant Constructs and Methods of Use**

Of particular interest is the use of the nucleotide sequences in recombinant  
25 DNA constructs to direct the transcription or transcription and translation (expression) of the prenyltransferase or tocopherol cyclase sequences of the present invention in a host plant cell. The expression constructs generally comprise a promoter functional in a host plant cell operably linked to a nucleic acid sequence encoding a

prenyltransferase or tocopherol cyclase of the present invention and a transcriptional termination region functional in a host plant cell.

A first nucleic acid sequence is "operably linked" or "operably associated" with a second nucleic acid sequence when the sequences are so arranged that the first  
5 nucleic acid sequence affects the function of the second nucleic acid sequence. Preferably, the two sequences are part of a single contiguous nucleic acid molecule and more preferably are adjacent. For example, a promoter is operably linked to a gene if the promoter regulates or mediates transcription of the gene in a cell.

Those skilled in the art will recognize that there are a number of promoters  
10 which are functional in plant cells, and have been described in the literature. Chloroplast and plastid specific promoters, chloroplast or plastid functional promoters, and chloroplast or plastid operable promoters are also envisioned.

One set of plant functional promoters are constitutive promoters such as the CaMV35S or FMV35S promoters that yield high levels of expression in most plant  
15 organs. Enhanced or duplicated versions of the CaMV35S and FMV35S promoters are useful in the practice of this invention (Odell, *et al.* (1985) *Nature* 313:810-812; Rogers, U.S. Patent Number 5,378, 619). In addition, it may also be preferred to bring about expression of the prenyltransferase or tocopherol cyclase gene in specific tissues of the plant, such as leaf, stem, root, tuber, seed, fruit, etc., and the promoter chosen  
20 should have the desired tissue and developmental specificity.

Of particular interest is the expression of the nucleic acid sequences of the present invention from transcription initiation regions which are preferentially expressed in a plant seed tissue. Examples of such seed preferential transcription initiation sequences include those sequences derived from sequences encoding plant  
25 storage protein genes or from genes involved in fatty acid biosynthesis in oilseeds. Examples of such promoters include the 5' regulatory regions from such genes as napin (Kridl *et al.*, *Seed Sci. Res.* 1:209:219 (1991)), phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean  $\alpha'$  subunit of  $\beta$ -conglycinin (soy 7s, (Chen *et al.*, *Proc. Natl. Acad. Sci.*, 83:8560-8564 (1986))) and oleosin.

It may be advantageous to direct the localization of proteins conferring prenyltransferase or tocopherol cyclase to a particular subcellular compartment, for example, to the mitochondrion, endoplasmic reticulum, vacuoles, chloroplast or other plastidic compartment. For example, where the genes of interest of the present invention will be targeted to plastids, such as chloroplasts, for expression, the constructs will also employ the use of sequences to direct the gene to the plastid. Such sequences are referred to herein as chloroplast transit peptides (CTP) or plastid transit peptides (PTP). In this manner, where the gene of interest is not directly inserted into the plastid, the expression construct will additionally contain a gene encoding a transit peptide to direct the gene of interest to the plastid. The chloroplast transit peptides may be derived from the gene of interest, or may be derived from a heterologous sequence having a CTP. Such transit peptides are known in the art. See, for example, Von Heijne *et al.* (1991) *Plant Mol. Biol. Rep.* 9:104-126; Clark *et al.* (1989) *J. Biol. Chem.* 264:17544-17550; della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968; Romer *et al.* (1993) *Biochem. Biophys. Res Commun.* 196:1414-1421; and, Shah *et al.* (1986) *Science* 233:478-481.

Depending upon the intended use, the constructs may contain the nucleic acid sequence which encodes the entire prenyltransferase or tocopherol cyclase protein, or a portion thereof. For example, where antisense inhibition of a given prenyltransferase or tocopherol cyclase protein is desired, the entire prenyltransferase or tocopherol cyclase sequence is not required. Furthermore, where prenyltransferase or tocopherol cyclase sequences used in constructs are intended for use as probes, it may be advantageous to prepare constructs containing only a particular portion of a prenyltransferase or tocopherol cyclase encoding sequence, for example a sequence which is discovered to encode a highly conserved prenyltransferase or tocopherol cyclase region.

The skilled artisan will recognize that there are various methods for the inhibition of expression of endogenous sequences in a host cell. Such methods include, but are not limited to, antisense suppression (Smith, *et al.* (1988) *Nature*

334:724-726), co-suppression (Napoli, *et al.* (1989) *Plant Cell* 2:279-289), ribozymes (PCT Publication WO 97/10328), and combinations of sense and antisense Waterhouse, *et al.* (1998) *Proc. Natl. Acad. Sci. USA* 95:13959-13964. Methods for the suppression of endogenous sequences in a host cell typically employ the  
5 transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well as non-coding regions of the endogenous sequence.

Regulatory transcript termination regions may be provided in plant expression constructs of this invention as well. Transcript termination regions may be provided  
10 by the DNA sequence encoding the prenyltransferase or tocopherol cyclase or a convenient transcription termination region derived from a different gene source, for example, the transcript termination region which is naturally associated with the transcript initiation region. The skilled artisan will recognize that any convenient transcript termination region which is capable of terminating transcription in a plant  
15 cell may be employed in the constructs of the present invention.

Alternatively, constructs may be prepared to direct the expression of the prenyltransferase or tocopherol cyclase sequences directly from the host plant cell plastid. Such constructs and methods are known in the art and are generally described, for example, in Svab, *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-  
20 8530 and Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917 and in U.S. Patent Number 5,693,507.

The prenyltransferase or tocopherol cyclase constructs of the present invention can be used in transformation methods with additional constructs providing for the expression of other nucleic acid sequences encoding proteins involved in the  
25 production of tocopherols, or tocopherol precursors such as homogentisic acid and/or phytylpyrophosphate. Nucleic acid sequences encoding proteins involved in the production of homogentisic acid are known in the art, and include but not are limited to, 4-hydroxyphenylpyruvate dioxygenase (HPPD, EC 1.13.11.27) described for example, by Garcia, *et al.* ((1999) *Plant Physiol.* 119(4):1507-1516), mono or

bifunctional *tyrA* (described for example by Xia, *et al.* (1992) *J. Gen. Microbiol.* 138:1309-1316, and Hudson, *et al.* (1984) *J. Mol. Biol.* 180:1023-1051), Oxygenase, 4-hydroxyphenylpyruvate di- (9CI), 4-Hydroxyphenylpyruvate dioxygenase; p-Hydroxyphenylpyruvate dioxygenase; p-Hydroxyphenylpyruvate hydroxylase; 5 p-Hydroxyphenylpyruvate oxidase; p-Hydroxyphenylpyruvic acid hydroxylase; p-Hydroxyphenylpyruvic hydroxylase; p-Hydroxyphenylpyruvic oxidase), 4-hydroxyphenylacetate, NAD(P)H:oxygen oxidoreductase (1-hydroxylating); 4-hydroxyphenylacetate 1-monooxygenase, and the like. In addition, constructs for the expression of nucleic acid sequences encoding proteins involved in the production of phytylpyrophosphate can also be employed with the prenyltransferase or tocopherol 10 cyclase constructs of the present invention. Nucleic acid sequences encoding proteins involved in the production of phytylpyrophosphate are known in the art, and include, but are not limited to geranylgeranylpyrophosphate synthase (GGPPS), geranylgeranylpyrophosphate reductase (GGH), 1-deoxyxylulose-5-phosphate 15 synthase, 1- deoxy-D-xylulose-5-phosphate reductoisomerase, 4-diphosphocytidyl-2-C-methylerythritol synthase, isopentyl pyrophosphate isomerase.

The prenyltransferase or tocopherol cyclase sequences of the present invention find use in the preparation of transformation constructs having a second expression cassette for the expression of additional sequences involved in tocopherol 20 biosynthesis. Additional tocopherol biosynthesis sequences of interest in the present invention include, but are not limited to gamma-tocopherol methyltransferase (Shintani, *et al.* (1998) *Science* 282(5396):2098-2100), tocopherol cyclase, and tocopherol methyltransferase.

A plant cell, tissue, organ, or plant into which the recombinant DNA 25 constructs containing the expression constructs have been introduced is considered transformed, transfected, or transgenic. A transgenic or transformed cell or plant also includes progeny of the cell or plant and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered

phenotype resulting from the presence of a prenyltransferase or tocopherol cyclase nucleic acid sequence.

Plant expression or transcription constructs having a prenyltransferase or tocopherol cyclase as the DNA sequence of interest for increased or decreased  
5 expression thereof may be employed with a wide variety of plant life, particularly, plant life involved in the production of vegetable oils for edible and industrial uses. Particularly preferred plants for use in the methods of the present invention include, but are not limited to: *Acacia*, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, broccoli,  
10 brussels sprouts, cabbage, canola, cantaloupe, carrot, cassava, cauliflower, celery, cherry, chicory, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, garlic, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom, nectarine, nut, oat, oil palm, oil seed rape, okra, onion, orange, an  
15 ornamental plant, papaya, parsley, pea, peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, triticale, turf, turnip, a vine, watermelon, wheat, yams,  
20 and zucchini.

Most especially preferred are temperate oilseed crops. Temperate oilseed crops of interest include, but are not limited to, rapeseed (Canola and High Erucic Acid varieties), sunflower, safflower, cotton, soybean, peanut, coconut and oil palms, and corn. Depending on the method for introducing the recombinant constructs into  
25 the host cell, other DNA sequences may be required. Importantly, this invention is applicable to dicotyledons and monocotyledons species alike and will be readily applicable to new and/or improved transformation and regulation techniques.

Of particular interest, is the use of prenyltransferase or tocopherol cyclase constructs in plants to produce plants or plant parts, including, but not limited to

leaves, stems, roots, reproductive, and seed, with a modified content of tocopherols in plant parts having transformed plant cells.

For immunological screening, antibodies to the protein can be prepared by injecting rabbits or mice with the purified protein or portion thereof, such methods of preparing antibodies being well known to those in the art. Either monoclonal or polyclonal antibodies can be produced, although typically polyclonal antibodies are more useful for gene isolation. Western analysis may be conducted to determine that a related protein is present in a crude extract of the desired plant species, as determined by cross-reaction with the antibodies to the encoded proteins. When cross-reactivity is observed, genes encoding the related proteins are isolated by screening expression libraries representing the desired plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory, Cold Spring Harbor, New York).

To confirm the activity and specificity of the proteins encoded by the identified nucleic acid sequences as prenyltransferase or tocopherol cyclase enzymes, *in vitro* assays are performed in insect cell cultures using baculovirus expression systems. Such baculovirus expression systems are known in the art and are described by Lee, *et al.* U.S. Patent Number 5,348,886, the entirety of which is herein incorporated by reference.

In addition, other expression constructs may be prepared to assay for protein activity utilizing different expression systems. Such expression constructs are transformed into yeast or prokaryotic host and assayed for prenyltransferase or tocopherol cyclase activity. Such expression systems are known in the art and are readily available through commercial sources.

In addition to the sequences described in the present invention, DNA coding sequences useful in the present invention can be derived from algae, fungi, bacteria, mammalian sources, plants, etc. Homology searches in existing databases using



signature sequences corresponding to conserved nucleotide and amino acid sequences of prenyltransferase or tocopherol cyclase can be employed to isolate equivalent, related genes from other sources such as plants and microorganisms. Searches in EST databases can also be employed. Furthermore, the use of DNA sequences  
5 encoding enzymes functionally enzymatically equivalent to those disclosed herein, wherein such DNA sequences are degenerate equivalents of the nucleic acid sequences disclosed herein in accordance with the degeneracy of the genetic code, is also encompassed by the present invention. Demonstration of the functionality of coding sequences identified by any of these methods can be carried out by  
10 complementation of mutants of appropriate organisms, such as *Synechocystis*, *Shewanella*, yeast, *Pseudomonas*, *Rhodobacteria*, etc., that lack specific biochemical reactions, or that have been mutated. The sequences of the DNA coding regions can be optimized by gene resynthesis, based on codon usage, for maximum expression in particular hosts.

15 For the alteration of tocopherol production in a host cell, a second expression construct can be used in accordance with the present invention. For example, the prenyltransferase or tocopherol cyclase expression construct can be introduced into a host cell in conjunction with a second expression construct having a nucleotide sequence for a protein involved in tocopherol biosynthesis.

20 The method of transformation in obtaining such transgenic plants is not critical to the instant invention, and various methods of plant transformation are currently available. Furthermore, as newer methods become available to transform crops, they may also be directly applied hereunder. For example, many plant species naturally susceptible to *Agrobacterium* infection may be successfully transformed via tripartite  
25 or binary vector methods of *Agrobacterium* mediated transformation. In many instances, it will be desirable to have the construct bordered on one or both sides by T-DNA, particularly having the left and right borders, more particularly the right border. This is particularly useful when the construct uses *A. tumefaciens* or *A. rhizogenes* as a mode for transformation, although the T-DNA borders may find use

with other modes of transformation. In addition, techniques of microinjection, DNA particle bombardment, and electroporation have been developed which allow for the transformation of various monocot and dicot plant species.

Normally, included with the DNA construct will be a structural gene having  
5 the necessary regulatory regions for expression in a host and providing for selection of transformant cells. The gene may provide for resistance to a cytotoxic agent, e.g. antibiotic, heavy metal, toxin, etc., complementation providing prototrophy to an auxotrophic host, viral immunity or the like. Depending upon the number of different host species the expression construct or components thereof are introduced, one or  
10 more markers may be employed, where different conditions for selection are used for the different hosts.

Where *Agrobacterium* is used for plant cell transformation, a vector may be used which may be introduced into the *Agrobacterium* host for homologous recombination with T-DNA or the Ti- or Ri-plasmid present in the *Agrobacterium*  
15 host. The Ti- or Ri-plasmid containing the T-DNA for recombination may be armed (capable of causing gall formation) or disarmed (incapable of causing gall formation), the latter being permissible, so long as the *vir* genes are present in the transformed *Agrobacterium* host. The armed plasmid can give a mixture of normal plant cells and gall.

20 In some instances where *Agrobacterium* is used as the vehicle for transforming host plant cells, the expression or transcription construct bordered by the T-DNA border region(s) will be inserted into a broad host range vector capable of replication in *E. coli* and *Agrobacterium*, there being broad host range vectors described in the literature. Commonly used is pRK2 or derivatives thereof. See, for example, Ditta, *et al.*, (Proc. Nat. Acad. Sci., U.S.A. (1980) 77:7347-7351) and EPA 0 120 515, which  
25 are incorporated herein by reference. Alternatively, one may insert the sequences to be expressed in plant cells into a vector containing separate replication sequences, one of which stabilizes the vector in *E. coli*, and the other in *Agrobacterium*. See, for example, McBride, *et al.* (Plant Mol. Biol. (1990) 14:269-276), wherein the pRIHRI

(Jouanin, *et al.*, *Mol. Gen. Genet.* (1985) 201:370-374) origin of replication is utilized and provides for added stability of the plant expression vectors in host *Agrobacterium* cells.

Included with the expression construct and the T-DNA will be one or more  
5 markers, which allow for selection of transformed *Agrobacterium* and transformed plant cells. A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, kanamycin, the aminoglycoside G418, hygromycin, or the like. The particular marker employed is not essential to this invention, one or another marker  
10 being preferred depending on the particular host and the manner of construction.

For transformation of plant cells using *Agrobacterium*, explants may be combined and incubated with the transformed *Agrobacterium* for sufficient time for transformation, the bacteria killed, and the plant cells cultured in an appropriate selective medium. Once callus forms, shoot formation can be encouraged by  
15 employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be grown to seed and the seed used to establish repetitive generations and for isolation of vegetable oils.

There are several possible ways to obtain the plant cells of this invention  
20 which contain multiple expression constructs. Any means for producing a plant comprising a construct having a DNA sequence encoding the expression construct of the present invention, and at least one other construct having another DNA sequence encoding an enzyme are encompassed by the present invention. For example, the expression construct can be used to transform a plant at the same time as the second  
25 construct either by inclusion of both expression constructs in a single transformation vector or by using separate vectors, each of which express desired genes. The second construct can be introduced into a plant which has already been transformed with the prenyltransferase or tocopherol cyclase expression construct, or alternatively, transformed plants, one expressing the prenyltransferase or tocopherol cyclase

construct and one expressing the second construct, can be crossed to bring the constructs together in the same plant.

Transgenic plants of the present invention may be produced from tissue culture, and subsequent generations grown from seed. Alternatively, transgenic plants  
5 may be grown using apomixis. Apomixis is a genetically controlled method of reproduction in plants where the embryo is formed without union of an egg and a sperm. There are three basic types of apomictic reproduction: 1) apospory where the embryo develops from a chromosomally unreduced egg in an embryo sac derived from the nucleus, 2) diplospory where the embryo develops from an unreduced egg in  
10 an embryo sac derived from the megaspore mother cell, and 3) adventitious embryony where the embryo develops directly from a somatic cell. In most forms of apomixis, pseudogamy or fertilization of the polar nuclei to produce endosperm is necessary for seed viability. In apospory, a nurse cultivar can be used as a pollen source for endosperm formation in seeds. The nurse cultivar does not affect the genetics of the  
15 aposporous apomictic cultivar since the unreduced egg of the cultivar develops parthenogenetically, but makes possible endosperm production. Apomixis is economically important, especially in transgenic plants, because it causes any genotype, no matter how heterozygous, to breed true. Thus, with apomictic reproduction, heterozygous transgenic plants can maintain their genetic fidelity  
20 throughout repeated life cycles. Methods for the production of apomictic plants are known in the art. See, U.S. Patent No. 5,811,636, which is herein incorporated by reference in its entirety.

The nucleic acid sequences of the present invention can be used in constructs to provide for the expression of the sequence in a variety of host cells, both  
25 prokaryotic eukaryotic. Host cells of the present invention preferably include monocotyledenous and dicotyledenous plant cells.

In general, the skilled artisan is familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of macromolecules (e.g., DNA molecules, plasmids, etc.), generation of

recombinant organisms and the screening and isolating of clones, (see for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989); Maliga *et al.*, *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995), the entirety of which is herein incorporated by reference; Birren *et al.*,  
5 *Genome Analysis: Analyzing DNA*, 1, Cold Spring Harbor, New York, the entirety of which is herein incorporated by reference).

Methods for the expression of sequences in insect host cells are known in the art. Baculovirus expression vectors are recombinant insect viruses in which the coding sequence for a chosen foreign gene has been inserted behind a baculovirus promoter  
10 in place of the viral gene, e.g., polyhedrin (Smith and Summers, U.S. Pat. No., 4,745,051, the entirety of which is incorporated herein by reference). Baculovirus expression vectors are known in the art, and are described for example in Doerfler, *Curr. Top. Microbiol. Immunol.* 131:51-68 (1968); Luckow and Summers, *Bio/Technology* 6:47-55 (1988a); Miller, *Annual Review of Microbiol.* 42:177-199  
15 (1988); Summers, *Curr. Comm. Molecular Biology*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988); Summers and Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Ag. Exper. Station Bulletin No. 1555 (1988), the entireties of which is herein incorporated by reference)

Methods for the expression of a nucleic acid sequence of interest in a fungal  
20 host cell are known in the art. The fungal host cell may, for example, be a yeast cell or a filamentous fungal cell. Methods for the expression of DNA sequences of interest in yeast cells are generally described in "Guide to yeast genetics and molecular biology", Guthrie and Fink, eds. *Methods in enzymology*, Academic Press, Inc. Vol 194 (1991) and *Gene expression technology*", Goeddel ed, *Methods in Enzymology*, Academic  
25 Press, Inc., Vol 185 (1991).

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC, Manassas, VA), such as HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells and a number of other cell lines.

Suitable promoters for mammalian cells are also known in the art and include, but are not limited to, viral promoters such as that from Simian Virus 40 (SV40) (Fiers *et al.*, *Nature* 273:113 (1978), the entirety of which is herein incorporated by reference), Rous sarcoma virus (RSV), adenovirus (ADV) and bovine papilloma virus (BPV).

- 5 Mammalian cells may also require terminator sequences and poly-A addition sequences. Enhancer sequences which increase expression may also be included and sequences which promote amplification of the gene may also be desirable (for example methotrexate resistance genes).

- Vectors suitable for replication in mammalian cells are well known in the art,  
10 and may include viral replicons, or sequences which insure integration of the appropriate sequences encoding epitopes into the host genome. Plasmid vectors that greatly facilitate the construction of recombinant viruses have been described (*see*, for example, Mackett *et al.*, *J Virol.* 49:857 (1984); Chakrabarti *et al.*, *Mol. Cell. Biol.* 5:3403 (1985); Moss, In: *Gene Transfer Vectors For Mammalian Cells* (Miller and  
15 Calos, eds., Cold Spring Harbor Laboratory, N.Y., p. 10, (1987); all of which are herein incorporated by reference in their entirety).

- The invention also includes plants and plant parts, such as seed, oil and meal derived from seed, and feed and food products processed from plants, which are enriched in tocopherols. Of particular interest is seed oil obtained from transgenic  
20 plants where the tocopherol level has been increased as compared to seed oil of a non-transgenic plant.

- The harvested plant material may be subjected to additional processing to further enrich the tocopherol content. The skilled artisan will recognize that there are many such processes or methods for refining, bleaching and degumming oil. United  
25 States Patent Number 5,932,261, issued August 3, 1999, discloses on such process, for the production of a natural carotene rich refined and deodorised oil by subjecting the oil to a pressure of less than 0.060 mbar and to a temperature of less than 200.degree. C. Oil distilled by this process has reduced free fatty acids, yielding a refined, deodorised oil where Vitamin E contained in the feed oil is substantially

retained in the processed oil. The teachings of this patent are incorporated herein by reference.

The invention now being generally described, it will be more readily  
5 understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

### EXAMPLES

#### Example 1: Identification of Prenyltransferase or tocopherol cyclase Sequences

10 PSI-BLAST (Altschul, *et al.* (1997) *Nuc Acid Res* 25:3389-3402) profiles were generated for both the straight chain and aromatic classes of prenyltransferases. To generate the straight chain profile, a prenyl- transferase from *Porphyr*  
a *purpurea* (Genbank accession 1709766) was used as a query against the NCBI non-redundant protein database. The *E. coli* enzyme involved in the formation of ubiquinone, ubiA  
15 (genbank accession 1790473) was used as a starting sequence to generate the aromatic prenyltransferase profile. These profiles were used to search public and proprietary DNA and protein data bases. In *Arabidopsis* six putative prenyltransferases of the straight-chain class were identified, ATPT1, (SEQ ID NO:9), ATPT7 (SEQ ID  
NO:10), ATPT8 (SEQ ID NO:11), ATPT9 (SEQ ID NO:13), ATPT10 (SEQ ID  
20 NO:14), and ATPT11 (SEQ ID NO:15), and six were identified of the aromatic class, ATPT2 (SEQ ID NO:1), ATPT3 (SEQ ID NO:3), ATPT4 (SEQ ID NO:5), ATPT5 (SEQ ID NO:7), ATPT6 (SEQ ID NO:8), and ATPT12 (SEQ ID NO:16). Additional prenyltransferase sequences from other plants related to the aromatic class of prenyltransferases, such as soy (SEQ ID NOs: 19-23, the deduced amino acid  
25 sequence of SEQ ID NO:23 is provided in SEQ ID NO:24) and maize (SEQ ID NOs:25-29, and 31) are also identified. The deduced amino acid sequence of ZMPT5 (SEQ ID NO:29) is provided in SEQ ID NO:30.

Searches are performed on a Silicon Graphics Unix computer using additional Bioaccelerator hardware and GenWeb software supplied by Compugen Ltd. This

software and hardware enables the use of the Smith-Waterman algorithm in searching DNA and protein databases using profiles as queries. The program used to query protein databases is profilesearch. This is a search where the query is not a single sequence but a profile based on a multiple alignment of amino acid or nucleic acid sequences. The profile is used to query a sequence data set, i.e., a sequence database. The profile contains all the pertinent information for scoring each position in a sequence, in effect replacing the "scoring matrix" used for the standard query searches. The program used to query nucleotide databases with a protein profile is tprofilesearch. Tprofilesearch searches nucleic acid databases using an amino acid profile query. As the search is running, sequences in the database are translated to amino acid sequences in six reading frames. The output file for tprofilesearch is identical to the output file for profilesearch except for an additional column that indicates the frame in which the best alignment occurred.

The Smith-Waterman algorithm, (Smith and Waterman (1981) *supra*), is used to search for similarities between one sequence from the query and a group of sequences contained in the database. E score values as well as other sequence information, such as conserved peptide sequences are used to identify related sequences.

To obtain the entire coding region corresponding to the *Arabidopsis* prenyltransferase sequences, synthetic oligo-nucleotide primers are designed to amplify the 5' and 3' ends of partial cDNA clones containing prenyltransferase sequences. Primers are designed according to the respective *Arabidopsis* prenyltransferase sequences and used in Rapid Amplification of cDNA Ends (RACE) reactions (Frohman *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) using the Marathon cDNA amplification kit (Clontech Laboratories Inc, Palo Alto, CA).

Amino acid sequence alignments between ATPT2 (SEQ ID NO:2), ATPT3 (SEQ ID NO:4), ATPT4 (SEQ ID NO:6), ATPT8 (SEQ ID NO:12), and ATPT12 (SEQ ID NO:17) are performed using ClustalW (Figure 1), and the percent identity and similarities are provided in Table 1 below.



Table 1:

	ATPT2	ATPT3	ATPT4	ATPT8	ATPT12
ATPT2 % Identity	12	13	11	15	
% similar	25	25	22	32	
% Gap	17	20	20	9	
ATPT3 % Identity		12	6	22	
% similar		29	16	38	
% Gap		20	24	14	
ATPT4 % Identity			9	14	
% similar			18	29	
% Gap			26	19	
ATPT8 % Identity				7	
% similar				19	
% Gap				20	
ATPT12 % Identity					
% similar					
% Gap					

**Example 2: Preparation of Prenyl Transferase Expression Constructs**

- 5           A plasmid containing the napin cassette derived from pCGN3223 (described in USPN 5,639,790, the entirety of which is incorporated herein by reference) was modified to make it more useful for cloning large DNA fragments containing multiple restriction sites, and to allow the cloning of multiple napin fusion genes into plant binary transformation vectors. An adapter comprised of the self annealed
- 10   oligonucleotide of sequence
- CGCGATTAAATGGCGCGCCCTGCAGGCGCCGCTGCAGGGCGCGCCAT
- TAAAT (SEQ ID NO:40) was ligated into the cloning vector pBC SK+ (Stratagene) after digestion with the restriction endonuclease BssHII to construct vector

pCGN7765. Plasmids pCGN3223 and pCGN7765 were digested with NotI and ligated together. The resultant vector, pCGN7770, contains the pCGN7765 backbone with the napin seed specific expression cassette from pCGN3223.

The cloning cassette, pCGN7787, essentially the same regulatory elements as pCGN7770, with the exception of the napin regulatory regions of pCGN7770 have been replaced with the double CAMV 35S promoter and the trnI polyadenylation and transcriptional termination region.

A binary vector for plant transformation, pCGN5139, was constructed from pCGN1558 (McBride and Summerfelt, (1990) Plant Molecular Biology, 14:269-276). The polylinker of pCGN1558 was replaced as a HindIII/Asp718 fragment with a polylinker containing unique restriction endonuclease sites, AscI, PacI, XbaI, SmaI, BamHI, and NotI. The Asp718 and HindIII restriction endonuclease sites are retained in pCGN5139.

A series of turbo binary vectors are constructed to allow for the rapid cloning of DNA sequences into binary vectors containing transcriptional initiation regions (promoters) and transcriptional termination regions.

The plasmid pCGN8618 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:41) and 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:42) into SalI/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was excised from pCGN8618 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8622.

The plasmid pCGN8619 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC -3' (SEQ ID NO:43) and 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:44) into SalI/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was removed from pCGN8619 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8623.

The plasmid pCGN8620 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGGAGCT -3' (SEQ ID NO:45) and 5'-CCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:46) into SalI/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8620 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8624.

The plasmid pCGN8621 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCCAGCT -3' (SEQ ID NO:47) and 5'-GGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:48) into

SalI/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8621 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8625.

The plasmid construct pCGN8640 is a modification of pCGN8624 described above. A 938bp PstI fragment isolated from transposon Tn7 which encodes bacterial spectinomycin and streptomycin resistance (Fling et al. (1985), *Nucleic Acids Research* 13(19):7095-7106), a determinant for *E. coli* and *Agrobacterium* selection, was blunt ended with Pfu polymerase. The blunt ended fragment was ligated into pCGN8624 that had been digested with SpeI and blunt ended with Pfu polymerase. The region containing the PstI fragment was sequenced to confirm both the insert orientation and the integrity of cloning junctions.

The spectinomycin resistance marker was introduced into pCGN8622 and pCGN8623 as follows. A 7.7 Kbp AvrII-SnaBI fragment from pCGN8640 was ligated to a 10.9 Kbp AvrII-SnaBI fragment from pCGN8623 or pCGN8622, described above. The resulting plasmids were pCGN8641 and pCGN8643, respectively.

The plasmid pCGN8644 was constructed by ligating oligonucleotides 5'-GATCACCTGCAGGAAGCTTGC GGCCGCGGATCCAATGCA-3' (SEQ ID NO:49) and 5'-TTGGATCCGCGGCCGCAAGCTTCCTGCAGGT-3' (SEQ ID NO:50) into BamHI-PstI digested pCGN8640.

Synthetic oligonucleotides were designed for use in Polymerase Chain Reactions (PCR) to amplify the coding sequences of ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 for the preparation of expression constructs and are provided in Table 2 below.

5 **Table 2:**

Name	Restriction Site	Sequence	SEQ ID NO:
ATPT2	5' NotI	GGATCCGCGGCCGCGACAATGGAGTC TCTGCTCTCTAGTTCT	51
ATPT2	3' SseI	GGATCCTGCAGGTCACCTCAAAAAA GGTAACAGCAAGT	52
ATPT3	5' NotI	GGATCCGCGGCCGCGACAATGGCGTT TTTGGGCTCTCCCGTGTTT	53
ATPT3	3' SseI	GGATCCTGCAGGTTATTGAAAACCTT CTTCCAAGTACAAC	54
ATPT4	5' NotI	GGATCCGCGGCCGCGACAATGTGGCG AAGATCTGTTGTT	55
ATPT4	3' SseI	GGATCCTGCAGGTCATGGAGAGTAG AAGGAAGGAGCT	56
ATPT8	5' NotI	GGATCCGCGGCCGCGACAATGGTACT TGCCGAGGTTCCAAAGCTTGCCCTCT	57
ATPT8	3' SseI	GGATCCTGCAGGTCACCTGTTTCTGG TGATGACTCTAT	58
ATPT12	5' NotI	GGATCCGCGGCCGCGACAATGACTTC GATTCTCAACACT	59
ATPT12	3' SseI	GGATCCTGCAGGTCAGTGTGCGAT GCTAATGCCGT	60

The coding sequences of ATPT2, ATPT3, ATPT4, ATPT8, and ATPT12 were all amplified using the respective PCR primers shown in Table 2 above and cloned into the TopoTA vector (Invitrogen). Constructs containing the respective prenyltransferase  
 10 sequences were digested with NotI and Sse8387I and cloned into the turbobinary vectors described above.

The sequence encoding ATPT2 prenyltransferase was cloned in the sense orientation into pCGN8640 to produce the plant transformation construct pCGN10800 (Figure 2). The ATPT2 sequence is under control of the 35S promoter.

The ATPT2 sequence was also cloned in the antisense orientation into the construct pCGN8641 to create pCGN10801 (Figure 3). This construct provides for the antisense expression of the ATPT2 sequence from the napin promoter.

The ATPT2 coding sequence was also cloned in the sense orientation into the  
5 vector pCGN8643 to create the plant transformation construct pCGN10822

The ATPT2 coding sequence was also cloned in the antisense orientation into the vector pCGN8644 to create the plant transformation construct pCGN10803 (Figure 4).

The ATPT4 coding sequence was cloned into the vector pCGN864 to create the plant transformation construct pCGN10806 (Figure 5). The ATPT2 coding sequence was  
10 cloned into the vector TopoTA™ vector from Invitrogen, to create the plant transformation construct pCGN10807 (Figure 6). The ATPT3 coding sequence was cloned into the TopoTA vector to create the plant transformation construct pCGN10808 (Figure 7). The ATPT3 coding sequence was cloned in the sense orientation into the vector pCGN8640 to create the plant transformation construct pCGN10809 (Figure 8). The  
15 ATPT3 coding sequence was cloned in the antisense orientation into the vector pCGN8641 to create the plant transformation construct pCGN10810 (Figure 9). The ATPT3 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10811 (Figure 10). The ATPT3 coding sequence was cloned into the vector pCGN8644 to create the plant transformation construct  
20 pCGN10812 (Figure 11). The ATPT4 coding sequence was cloned into the vector pCGN8640 to create the plant transformation construct pCGN10813 (Figure 12). The ATPT4 coding sequence was cloned into the vector pCGN8641 to create the plant transformation construct pCGN10814 (Figure 13). The ATPT4 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct  
25 pCGN10815 (Figure 14). The ATPT4 coding sequence was cloned in the antisense orientation into the vector pCGN8644 to create the plant transformation construct pCGN10816 (Figure 15). The ATPT8 coding sequence was cloned in the sense orientation into the vector pCGN8643 to create the plant transformation construct pCGN10819 (Figure 17). The ATPT12 coding sequence was cloned into the vector

pCGN8640 to create the plant transformation construct pCGN10824 (Figure 18). The ATPT12 coding sequence was cloned into the vector pCGN8643 to create the plant transformation construct pCGN10825 (Figure 19). The ATPT8 coding sequence was cloned into the vector pCGN8640 to create the plant transformation construct  
5 pCGN10826 (Figure 20).

**Example 3: Plant Transformation with Prenyl Transferase Constructs**

Transgenic *Brassica* plants are obtained by *Agrobacterium*-mediated transformation as described by Radke *et al.* (*Theor. Appl. Genet.* (1988) 75:685-694;  
10 *Plant Cell Reports* (1992) 11:499-505). Transgenic *Arabidopsis thaliana* plants may be obtained by *Agrobacterium*-mediated transformation as described by Valverkens *et al.*, (*Proc. Nat. Acad. Sci.* (1988) 85:5536-5540), or as described by Bent *et al.* ((1994), *Science* 265:1856-1860), or Bechtold *et al.* ((1993), *C.R.Acad.Sci, Life Sciences* 316:1194-1199). Other plant species may be similarly transformed using  
15 related techniques.

Alternatively, microprojectile bombardment methods, such as described by Klein *et al.* (*Bio/Technology* 10:286-291) may also be used to obtain nuclear transformed plants.

20 **Example 4: Identification of Additional Prenyltransferases**

Additional BLAST searches were performed using the ATPT2 sequence, a sequence in the class of aromatic prenyltransferases. ESTs, and in some case, full-length coding regions, were identified in proprietary DNA libraries.

Soy full-length homologs to ATPT2 were identified by a combination of  
25 BLAST (using ATPT2 protein sequence) and 5' RACE. Two homologs resulted (SEQ ID NO:95 and SEQ ID NO:96). Translated amino acid sequences are provided by SEQ ID NO:97 and SEQ ID NO:98.

A rice est ATPT2 homolog is shown in SEQ ID NO:99 (obtained from BLAST using the wheat ATPT2 homolog).

Other homolog sequences were obtained using ATPT2 and PSI-BLAST, including est sequences from wheat (SEQ ID NO:100), leek (SEQ ID NOs:101 and 102), canola (SEQ ID NO:103), corn (SEQ ID NOs:104, 105 and 106), cotton (SEQ ID NO:107) and tomato (SEQ ID NO:108).

- 5 A PSI-Blast profile generated using the *E. coli* ubiA (genbank accession 1790473) sequence was used to analyze the *Synechocystis* genome. This analysis identified 5 open reading frames (ORFs) in the *Synechocystis* genome that were potentially prenyltransferases; slr0926 (annotated as ubiA (4-hydroxybenzoate-octaprenyltransferase, SEQ ID NO:32), slr1899 (annotated as ctaB (cytochrome c  
10 oxidase folding protein, SEQ ID NO:33), slr0056 (annotated as g4 (chlorophyll synthase 33 kd subunit, SEQ ID NO:34), slr1518 (annotated as menA (menaquinone biosynthesis protein, SEQ ID NO:35), and slr1736 (annotated as a hypothetical protein of unknown function (SEQ ID NO:36).

#### 15 4A. *Synechocystis* Knock-outs

To determine the functionality of these ORFs and their involvement, if any, in the biosynthesis of tocopherols, knockouts constructs were made to disrupt the ORF identified in *Synechocystis*.

- Synthetic oligos were designed to amplify regions from the 5' (5'-  
20 TAATGTGTACATTGTCGGCCTC (17365') (SEQ ID NO:61) and 5'-  
GCAATGTAACATCAGAGATTTTGAGACACAACGTGGCTTTCCACAATTCC  
CCGCACCGTC (1736kanpr1)) (SEQ ID NO:62) and 3' (5'-  
AGGCTAATAAGCACAAATGGGA (17363') (SEQ ID NO:63) and 5'-  
GGTATGAGTCAGCAACACCTTCTTCACGAGGCAGACCTCAGC  
25 GGAATTGGTTTAGGTTATCCC (1736kanpr2)) (SEQ ID NO:64) ends of the  
slr1736 ORF. The 1736kanpr1 and 1736kanpr2 oligos contained 20 bp of homology to the slr1736 ORF with an additional 40 bp of sequence homology to the ends of the kanamycin resistance cassette. Separate PCR steps were completed with these oligos and the products were gel purified and combined with the kanamycin resistance gene



from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The combined fragments were allowed to assemble without oligos under the following conditions: 94°C for 1 min, 55°C for 1 min, 72°C for 1 min plus 5 seconds per cycle for 40 cycles using pfu polymerase in 100ul reaction volume (Zhao, H and Arnold (1997) *Nucleic Acids Res.* 25(6):1307-1308). One microliter or five microliters of this assembly reaction was then amplified using 5' and 3' oligos nested within the ends of the ORF fragment, so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21681 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers. The *ubiA* 5' sequence was amplified using the primers 5'-GGATCCATGGTT GCCCAAACCCCATC (SEQ ID NO:65) and 5'-GCAATGTAACATCAGAGA TTTTGAGACACAACG TGGCTTTGGGTAAGCAACAATGACCGGC (SEQ ID NO:66). The 3' region was amplified using the synthetic oligonucleotide primers 5'-GAATTCTCAAAGCCAGCCCAGTAAC (SEQ ID NO:67) and 5'-GGTATGAGTC AGCAACACCTTCTTCACGAGGCAGACCTCAGCGGGTGCAGAAAAGGGTTTT CCC (SEQ ID NO:68). The amplification products were combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of the ORF fragment (5'-CCAGTGGTTTtaggctgtgtggc (SEQ ID NO:69) and 5'-CTGAGTTGGATGTATTGGATC (SEQ ID NO:70)), so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked

out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21682 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers. The sl11899 5' sequence was amplified using the primers 5'-GGATCCATGGTTACTT CGACAAAAATCC (SEQ ID NO:71) and 5'-GCAATGTAACATCAGAG ATTTTGAGACACAACGTGGCTTTGCTAGGCAACCGCTTAGTAC (SEQ ID NO:72). The 3' region was amplified using the synthetic oligonucleotide primers 5'-GAATTCTTAACCCAACAGTAAAGTTCCC (SEQ ID NO:73) and 5'-GGTATGAGTCAGC AACACCTTCTTCACGAGGCAGACCTCAGCGCCGGCATTGTCTTTTACATG (SEQ ID NO:74). The amplification products were combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of the ORF fragment (5'-GGAACCCTTGCAGCCGCTTC (SEQ ID NO:75) and 5'-GTATGCCCAACTGGTGCAGAGG (SEQ ID NO:76)), so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21679 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout constructs for the other sequences using the same method as described above, with the following primers. The slr0056 5' sequence was amplified using the primers 5'-GGATCCATGTCTGACACACAAAATACCG (SEQ ID NO:77) and 5'-GCAATGTAACATCAGAGATTTTGAGACACAACGTGGCTTTCGCCAATACC AGCCACCAACAG (SEQ ID NO:78). The 3' region was amplified using the

synthetic oligonucleotide primers 5'- GAATTCTCAAAT CCCCGCATGGCCTAG (SEQ ID NO:79) and 5'- GGTATGAGTCAGCAACACCTTCTTCACGAGGCAGACCTCAGCGGCCTACG GCTTGGACGTGTGGG (SEQ ID NO:80). The amplification products were

5 combined with the kanamycin resistance gene from puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of the ORF fragment (5'- CACTTGGATTCCCCTGATCTG (SEQ ID NO:81) and 5'- GCAATACCCGCTTGGAAAACG (SEQ ID NO:82)), so that the resulting product

10 contained 100-200 bp of the 5' end of the *Synechocystis* gene to be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21677 and used for *Synechocystis* transformation.

Primers were also synthesized for the preparation of *Synechocystis* knockout

15 constructs for the other sequences using the same method as described above, with the following primers. The slr1518 5' sequence was amplified using the primers 5'- GGATCCATGACCGAAT CTTCGCCCCTAGC (SEQ ID NO:83) and 5'- GCAATGTAACATCAGAGATTTTGA GACACAACGTGGC TTTCAATCCTAGGTAGCCGAGGCG (SEQ ID NO:84). The 3' region was

20 amplified using the synthetic oligonucleotide primers 5'- GAATTCTTAGCCCAGGCC AGCCCAGCC (SEQ ID NO:85) and 5'- GGTATGAGTCAGCAACACCTTCTTCACGA GGCAGACCTCAGCGGGGAATTGATTGTGTTAATTACC (SEQ ID NO:86). The amplification products were combined with the kanamycin resistance gene from

25 puc4K (Pharmacia) that had been digested with *HincII* and gel purified away from the vector backbone. The annealed fragment was amplified using 5' and 3' oligos nested within the ends of the ORF fragment (5'- GCGATCGCCATTATCGCTTGG (SEQ ID NO:87) and 5'- GCAGACTGGCAATTATCAGTAACG (SEQ ID NO:88)), so that the resulting product contained 100-200 bp of the 5' end of the *Synechocystis* gene to

be knocked out, the kanamycin resistance cassette, and 100-200 bp of the 3' end of the gene to be knocked out. This PCR product was then cloned into the vector pGemT easy (Promega) to create the construct pMON21680 and used for *Synechocystis* transformation.

5

#### 4B. Transformation of *Synechocystis*

Cells of *Synechocystis* 6803 were grown to a density of approximately  $2 \times 10^8$  cells per ml and harvested by centrifugation. The cell pellet was re-suspended in fresh BG-11 medium (ATCC Medium 616) at a density of  $1 \times 10^9$  cells per ml and used  
10 immediately for transformation. One-hundred microliters of these cells were mixed with 5 ul of mini prep DNA and incubated with light at 30C for 4 hours. This mixture was then plated onto nylon filters resting on BG-11 agar supplemented with TES pH8 and allowed to grow for 12-18 hours. The filters were then transferred to BG-11 agar + TES + 5ug/ml kanamycin and allowed to grow until colonies appeared within 7-10  
15 days (Packer and Glazer, 1988). Colonies were then picked into BG-11 liquid media containing 5 ug/ml kanamycin and allowed to grow for 5 days. These cells were then transferred to Bg-11 media containing 10ug/ml kanamycin and allowed to grow for 5 days and then transferred to Bg-11 + kanamycin at 25ug/ml and allowed to grow for 5 days. Cells were then harvested for PCR analysis to determine the presence of a  
20 disrupted ORF and also for HPLC analysis to determine if the disruption had any effect on tocopherol levels.

PCR analysis of the *Synechocystis* isolates for slr1736 and sl11899 showed complete segregation of the mutant genome, meaning no copies of the wild type genome could be detected in these strains. This suggests that function of the native  
25 gene is not essential for cell function. HPLC analysis of these same isolates showed that the sl11899 strain had no detectable reduction in tocopherol levels. However, the strain carrying the knockout for slr1736 produced no detectable levels of tocopherol.

The amino acid sequences for the *Synechocystis* knockouts are compared using ClustalW, and are provided in Table 3 below. Provided are the percent identities,

percent similarity, and the percent gap. The alignment of the sequences is provided in Figure 21.

**Table 3:**

	Slr1736	slr0926	sll1899	slr0056	slr1518
slr1736 %identity	14	12	18	11	
%similar	29	30	34	26	
%gap	8	7	10	5	
slr0926 %identity		20	19	14	
%similar		39	32	28	
%gap		7	9	4	
sll1899 %identity			17	13	
%similar			29	29	
%gap			12	9	
slr0056 %identity				15	
%similar				31	
%gap				8	
slr1518 %identity					
%similar					
%gap					

5

Amino acid sequence comparisons are performed using various *Arabidopsis* prenyltransferase sequences and the *Synechocystis* sequences. The comparisons are presented in Table 4 below. Provided are the percent identities, percent similarity, and the percent gap. The alignment of the sequences is provided in Figure 22.

**Table 4:**

[illegible]

#### 4C. Phytyl Prenyltransferase Enzyme Assays

[<sup>3</sup>H] Homogentisic acid in 0.1% H<sub>3</sub>PO<sub>4</sub> (specific radioactivity 40 Ci/mmol).

Phytyl pyrophosphate was synthesized as described by Joo, *et al.* (1973) *Can J.*

- 5 *Biochem.* 51:1527. 2-methyl-6-phytylquinol and 2,3-dimethyl-5-phytylquinol were synthesized as described by Soll, *et al.* (1980) *Phytochemistry* 19:215. Homogentisic acid,  $\alpha$ ,  $\beta$ ,  $\delta$ , and  $\gamma$ -tocopherol, and tocol, were purchased commercially.

- The wild-type strain of *Synechocystis* sp. PCC 6803 was grown in BG11 medium with bubbling air at 30°C under 50  $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  fluorescent light, and 70% relative  
10 humidity. The growth medium of slr1736 knock-out (potential PPT) strain of this organism was supplemented with 25  $\mu\text{g mL}^{-1}$  kanamycin. Cells were collected from 0.25 to 1 liter culture by centrifugation at 5000 *g* for 10 min and stored at -80°C.

- Total membranes were isolated according to Zak's procedures with some modifications (Zak, *et al.* (1999) *Eur J. Biochem* 261:311). Cells were broken on a  
15 French press. Before the French press treatment, the cells were incubated for 1 hour with lysozyme (0.5%, w/v) at 30 °C in a medium containing 7 mM EDTA, 5 mM NaCl and 10 mM Hepes-NaOH, pH 7.4. The spheroplasts were collected by centrifugation at 5000 *g* for 10 min and resuspended at 0.1 - 0.5 mg chlorophyll·mL<sup>-1</sup> in 20 mM potassium phosphate buffer, pH 7.8. Proper amount of protease inhibitor cocktail and  
20 DNAase I from Boehringer Mannheim were added to the solution. French press treatments were performed two to three times at 100 MPa. After breakage, the cell suspension was centrifuged for 10 min at 5000*g* to pellet unbroken cells, and this was followed by centrifugation at 100 000 *g* for 1 hour to collect total membranes. The final pellet was resuspended in a buffer containing 50 mM Tris-HCL and 4 mM MgCl<sub>2</sub>.

- 25 Chloroplast pellets were isolated from 250 g of spinach leaves obtained from local markets. Devined leaf sections were cut into grinding buffer (2 l /250 g leaves) containing 2 mM EDTA, 1 mM MgCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, 0.33 M sorbitol, 0.1% ascorbic acid, and 50 mM Hepes at pH 7.5. The leaves were homogenized for 3 sec three times in a 1-L blender, and filtered through 4 layers of miracloth. The supernatant was then

centrifuged at 5000g for 6 min. The chloroplast pellets were resuspended in small amount of grinding buffer (Douce, *et al* Methods in Chloroplast Molecular Biology, 239 (1982))

Chloroplasts in pellets can be broken in three ways. Chloroplast pellets were first  
5 aliquoted in 1 mg of chlorophyll per tube, centrifuged at 6000 rpm for 2 min in microcentrifuge, and grinding buffer was removed. Two hundred microliters of Triton X-100 buffer (0.1% Triton X-100, 50 mM Tris-HCl pH 7.6 and 4 mM MgCl<sub>2</sub>) or swelling buffer (10 mM Tris pH 7.6 and 4 mM MgCl<sub>2</sub>) was added to each tube and incubated for ½ hour at 4°C. Then the broken chloroplast pellets were used for the  
10 assay immediately. In addition, broken chloroplasts can also be obtained by freezing in liquid nitrogen and stored at -80°C for ½ hour, then used for the assay.

In some cases chloroplast pellets were further purified with 40%/ 80% percoll gradient to obtain intact chloroplasts. The intact chloroplasts were broken with swelling buffer, then either used for assay or further purified for envelope membranes  
15 with 20.5%/ 31.8% sucrose density gradient (Sol, *et al* (1980) *supra*). The membrane fractions were centrifuged at 100 000g for 40 min and resuspended in 50 mM Tris-HCl pH 7.6, 4 mM MgCl<sub>2</sub>.

Various amounts of [<sup>3</sup>H]HGA, 40 to 60 µM unlabelled HGA with specific activity in the range of 0.16 to 4 Ci/mmol were mixed with a proper amount of 1M Tris-  
20 NaOH pH 10 to adjust pH to 7.6. HGA was reduced for 4 min with a trace amount of solid NaBH<sub>4</sub>. In addition to HGA, standard incubation mixture (final vol 1 mL) contained 50 mM Tris-HCl, pH 7.6, 3-5 mM MgCl<sub>2</sub>, and 100 µM phytyl pyrophosphate. The reaction was initiated by addition of *Synechocystis* total membranes, spinach chloroplast pellets, spinach broken chloroplasts, or spinach  
25 envelope membranes. The enzyme reaction was carried out for 2 hour at 23°C or 30°C in the dark or light. The reaction is stopped by freezing with liquid nitrogen, and stored at -80°C or directly by extraction.

A constant amount of tocol was added to each assay mixture and reaction products were extracted with a 2 mL mixture of chloroform/methanol (1:2, v/v) to give a



monophasic solution. NaCl solution (2 mL; 0.9%) was added with vigorous shaking. This extraction procedure was repeated three times. The organic layer containing the prenylquinones was filtered through a 20 µm filter, evaporated under N<sub>2</sub> and then resuspended in 100 µL of ethanol.

5        The samples were mainly analyzed by Normal-Phase HPLC method (Isocratic 90% Hexane and 10% Methyl-t-butyl ether), and use a Zorbax silica column, 4.6 x 250 mm. The samples were also analyzed by Reversed-Phase HPLC method (Isocratic 0.1% H<sub>3</sub>PO<sub>4</sub> in MeOH), and use a Vydac 201HS54 C18 column; 4.6 x 250 mm coupled with an All-tech C18 guard column. The amount of products were calculated  
10       based on the substrate specific radioactivity, and adjusted according to the % recovery based on the amount of internal standard.

          The amount of chlorophyll was determined as described in Arnon (1949) *Plant Physiol.* 24:1. Amount of protein was determined by the Bradford method using gamma globulin as a standard (Bradford, (1976) *Anal. Biochem.* 72:248)

15        Results of the assay demonstrate that 2-Methyl-6-Phytylplastoquinone is not produced in the *Synechocystis* slr1736 knockout preparations. The results of the phytyl prenyltransferase enzyme activity assay for the slr1736 knock out are presented in Figure 23.

#### 20        4D. Complementation of the slr1736 knockout with ATPT2

          In order to determine whether ATPT2 could complement the knockout of slr1736 in *Synechocystis* 6803, a plasmid was constructed to express the ATPT2 sequence from the TAC promoter. A vector, plasmid psl1211, was obtained from the lab of Dr. Himadri Pakrasi of Washington University, and is based on the plasmid  
25        RSF1010 which is a broad host range plasmid (Ng W.-O., Zentella R., Wang, Y., Taylor J-S. A., Pakrasi, H.B. 2000. *phrA*, the major photoreactivating factor in the cyanobacterium *Synechocystis* sp. strain PCC 6803 codes for a cyclobutane pyrimidine dimer specific DNA photolyase. *Arch. Microbiol.* (in press)). The ATPT2 gene was isolated from the vector pCGN10817 by PCR using the following primers.

ATPT2<sub>nco.pr</sub> 5'-CCATGGATTTCGAGTAAAGTTGTCGC (SEQ ID NO:89);  
ATPT2<sub>ri.pr</sub> 5'-GAATTCACTTCAAAAAAGGTAACAG (SEQ ID NO:90). These  
primers will remove approximately 112 BP from the 5' end of the ATPT2 sequence,  
which is thought to be the chloroplast transit peptide. These primers will also add an  
5 NcoI site at the 5' end and an EcoRI site at the 3' end which can be used for sub-  
cloning into subsequent vectors. The PCR product from using these primers and  
pCGN10817 was ligated into pGEM T easy and the resulting vector pMON21689 was  
confirmed by sequencing using the m13forward and m13reverse primers. The  
NcoI/EcoRI fragment from pMON21689 was then ligated with the EagI/EcoRI and  
10 EagI/NcoI fragments from psl1211 resulting in pMON21690. The plasmid  
pMON21690 was introduced into the slr1736 *Synechocystis* 6803 KO strain via  
conjugation. Cells of sl906 (a helper strain) and DH10B cells containing  
pMON21690 were grown to log phase (O.D. 600= 0.4) and 1 ml was harvested by  
centrifugation. The cell pellets were washed twice with a sterile BG-11 solution and  
15 resuspended in 200 ul of BG-11. The following was mixed in a sterile eppendorf  
tube: 50 ul SL906, 50 ul DH10B cells containing pMON21690, and 100 ul of a fresh  
culture of the slr1736 *Synechocystis* 6803 KO strain (O.D. 730 = 0.2-0.4). The cell  
mixture was immediately transferred to a nitrocellulose filter resting on BG-11 and  
incubated for 24 hours at 30C and 2500 LUX(50 ue) of light. The filter was then  
20 transferred to BG-11 supplemented with 10ug/ml Gentamycin and incubated as above  
for ~5 days. When colonies appeared, they were picked and grown up in liquid BG-  
11 + Gentamycin 10 ug/ml. (Elhai, J. and Wolk, P. 1988. Conjugal transfer of DNA  
to Cyanobacteria. *Methods in Enzymology* 167, 747-54) The liquid cultures were then  
assayed for tocopherols by harvesting 1ml of culture by centrifugation, extracting with  
25 ethanol/pyrogallol, and HPLC separation. The slr1736 *Synechocystis* 6803 KO strain,  
did not contain any detectable tocopherols, while the slr1736 *Synechocystis* 6803 KO  
strain transformed with pmon21690 contained detectable alpha tocopherol. A  
*Synechocystis* 6803 strain transformed with psl1211(vector control) produced alpha  
tocopherol as well.

#### 4E: Additional Evidence of Prenyltransferase Activity

To test the hypothesis that slr1736 or ATPT2 are sufficient as single genes to obtain phytyl prenyltransferase activity, both genes were expressed in SF9 cells and in yeast. When either slr1736 or ATPT2 were expressed in insect cells (Table 5) or in yeast, phytyl prenyltransferase activity was detectable in membrane preparations, whereas membrane preparations of the yeast vector control, or membrane preparations of insect cells did not exhibit phytyl prenyltransferase activity.

10 **Table 5:** Phytyl prenyltransferase activity

Enzyme source	Enzyme activity [pmol/mg x h]
slr1736 expressed in SF9 cells	20
ATPT2 expressed in SF9 cells	6
SF9 cell control	< 0.05
<i>Synechocystis</i> 6803	0.25
Spinach chloroplasts	0.20

#### Example 5: Transgenic Plant Analysis

##### 15 5A. *Arabidopsis*

*Arabidopsis* plants transformed with constructs for the sense or antisense expression of the ATPT proteins were analyzed by High Pressure Liquid Chromatography (HPLC) for altered levels of total tocopherols, as well as altered levels of specific tocopherols (alpha, beta, gamma, and delta tocopherol).

20 Extracts of leaves and seeds were prepared for HPLC as follows. For seed extracts, 10 mg of seed was added to 1 g of microbeads (Biospec) in a sterile microfuge tube to which 500 ul 1% pyrogallol (Sigma Chem)/ethanol was added. The mixture was shaken for 3 minutes in a mini Beadbeater (Biospec) on "fast" speed.

The extract was filtered through a 0.2  $\mu$ m filter into an autosampler tube. The filtered extracts were then used in HPLC analysis described below.

Leaf extracts were prepared by mixing 30-50 mg of leaf tissue with 1 g microbeads and freezing in liquid nitrogen until extraction. For extraction, 500  $\mu$ l 1% pyrogallol in ethanol was added to the leaf/bead mixture and shaken for 1 minute on a  
5 Beadbeater (Biospec) on "fast" speed. The resulting mixture was centrifuged for 4 minutes at 14,000 rpm and filtered as described above prior to HPLC analysis.

HPLC was performed on a Zorbax silica HPLC column (4.6 mm X 250 mm) with a fluorescent detection, an excitation at 290 nm, an emission at 336 nm, and  
10 bandpass and slits. Solvent A was hexane and solvent B was methyl-t-butyl ether. The injection volume was 20  $\mu$ l, the flow rate was 1.5 ml/min, the run time was 12 min (40°C) using the gradient (Table 6):

Table 6:

15	<u>Time</u>	<u>Solvent A</u>	<u>Solvent B</u>
	0 min.	90%	10%
	10 min.	90%	10%
	11 min.	25%	75%
	12 min.	90%	10%

20

Tocopherol standards in 1% pyrogallol/ ethanol were also run for comparison (alpha tocopherol, gamma tocopherol, beta tocopherol, delta tocopherol, and tocopherol (tocol) (all from Matreya).

Standard curves for alpha, beta, delta, and gamma tocopherol were calculated  
25 using Chemstation software. The absolute amount of component x is: Absolute amount of x =  $\text{Response}_x \times \text{RF}_x \times \text{dilution factor}$  where  $\text{Response}_x$  is the area of peak x,  $\text{RF}_x$  is the response factor for component x ( $\text{Amount}_x / \text{Response}_x$ ) and the dilution factor is 500  $\mu$ l. The ng/mg tissue is found by: total ng component/mg plant tissue.

Results of the HPLC analysis of seed extracts of transgenic *Arabidopsis* lines containing pMON10822 for the expression of ATPT2 from the napin promoter are provided in Figure 24.

HPLC analysis results of segregating T2 *Arabidopsis* seed tissue expressing  
5 the ATPT2 sequence from the napin promoter (pCGN10822) demonstrates an increased level of tocopherols in the seed. Total tocopherol levels are increased as much as 50% over the total tocopherol levels of non-transformed (wild-type) *Arabidopsis* plants (Figure 25). Homozygous progeny from the top 3 lines (T3 seed) have up to a two-fold (100%) increase in total tocopherol levels over control  
10 *Arabidopsis* seed ( Figure 26.)

Furthermore, increases of particular tocopherols are also increased in transgenic *Arabidopsis* plants expressing the ATPT2 nucleic acid sequence from the napin promoter. Levels of delta tocopherol in these lines are increased greater than 3 fold over the delta tocopherol levels obtained from the seeds of wild type *Arabidopsis*  
15 lines. Levels of gamma tocopherol in transgenic *Arabidopsis* lines expressing the ATPT2 nucleic acid sequence are increased as much as about 60% over the levels obtained in the seeds of non-transgenic control lines. Furthermore, levels of alpha tocopherol are increased as much as 3 fold over those obtained from non-transgenic control lines.

20 Results of the HPLC analysis of seed extracts of transgenic *Arabidopsis* lines containing pCGN10803 for the expression of ATPT2 from the enhanced 35S promoter (antisense orientation ) are provided in Figure 25. Two lines were identified that have reduced total tocopherols, up to a ten-fold decrease observed in T3 seed compared to control *Arabidopsis* (Figure 27.)

25

#### 5B. Canola

Brassica napus, variety SP30021, was transformed with pCGN10822 (napin-ATPT2-napin 3', sense orientation) using *Agrobacterium tumefaciens*-mediated

transformation. Flowers of the R0 plants were tagged upon pollination and developing seed was collected at 35 and 45 days after pollination (DAP).

Developing seed was assayed for tocopherol levels, as described above for *Arabidopsis*. Line 10822-1 shows a 20% increase of total tocopherols, compared to the wild-type control, at 45 DAP. Figure 28 shows total tocopherol levels measured in developing canola seed.

**Example 6: Sequences to Tocopherol Cyclase**

**6A. Preparation of the *slr1737* Knockout**

The *Synechocystis* sp. 6803 *slr1737* knockout was constructed by the following method. The GPS™-1 Genome Priming System (New England Biolabs) was used to insert, by a Tn7 Transposase system, a Kanamycin resistance cassette into *slr1737*. A plasmid from a *Synechocystis* genomic library clone containing 652 base pairs of the targeted orf (*Synechocystis* genome base pairs 1324051 – 1324703; the predicted orf base pairs 1323672 – 1324763, as annotated by Cyanobase) was used as target DNA. The reaction was performed according to the manufacturers protocol. The reaction mixture was then transformed into *E. coli* DH10B electrocompetant cells and plated. Colonies from this transformation were then screened for transposon insertions into the target sequence by amplifying with M13 Forward and Reverse Universal primers, yielding a product of 652 base pairs plus ~1700 base pairs, the size of the transposon kanamycin cassette, for a total fragment size of ~2300 base pairs. After this determination, it was then necessary to determine the approximate location of the insertion within the targeted orf, as 100 base pairs of orf sequence was estimated as necessary for efficient homologous recombination in *Synechocystis*. This was accomplished through amplification reactions using either of the primers to the ends of the transposon, Primer S (5' end) or N (3' end), in combination with either a M13 Forward or Reverse primer. That is, four different primer combinations were used to map each potential knockout construct: Primer S – M13 Forward, Primer S – M13 Reverse, Primer N – M13 Forward, Primer N – M13 Reverse. The construct

used to transform *Synechocystis* and knockout slr1737 was determined to consist of a approximately 150 base pairs of slr1737 sequence on the 5' side of the transposon insertion and approximately 500 base pairs on the 3' side, with the transcription of the orf and kanamycin cassette in the same direction. The nucleic acid sequence of  
5 slr1737 is provided in SEQ ID NO:38 the deduced amino acid sequence is provided in SEQ ID NO:39.

Cells of *Synechocystis* 6803 were grown to a density of  $\sim 2 \times 10^8$  cells per ml and harvested by centrifugation. The cell pellet was re-suspended in fresh BG-11 medium at a density of  $1 \times 10^9$  cells per ml and used immediately for transformation.  
10 100 ul of these cells were mixed with 5 ul of mini prep DNA and incubated with light at 30C for 4 hours. This mixture was then plated onto nylon filters resting on BG-11 agar supplemented with TES pH8 and allowed to grow for 12-18 hours. The filters were then transferred to BG-11 agar + TES + 5ug/ml kanamycin and allowed to grow until colonies appeared within 7-10 days (Packer and Glazer, 1988). Colonies were  
15 then picked into BG-11 liquid media containing 5 ug/ml kanamycin and allowed to grow for 5 days. These cells were then transferred to Bg-11 media containing 10ug/ml kanamycin and allowed to grow for 5 days and then transferred to Bg-11 + kanamycin at 25ug/ml and allowed to grow for 5 days. Cells were then harvested for PCR analysis to determine the presence of a disrupted ORF and also for HPLC  
20 analysis to determine if the disruption had any effect on tocopherol levels.

PCR analysis of the *Synechocystis* isolates, using primers to the ends of the *slr1737* orf, showed complete segregation of the mutant genome, meaning no copies of the wild type genome could be detected in these strains. This suggests that function of the native gene is not essential for cell function. HPLC analysis of the  
25 strain carrying the knockout for *slr1737* produced no detectable levels of tocopherol.

#### 6B. The relation of slr1737 and slr1736

The slr1737 gene occurs in *Synechocystis* downstream and in the same orientation as slr1736, the phytol prenyltransferase. In bacteria this proximity often

indicates an operon structure and therefore an expression pattern that is linked in all genes belonging to this operon. Occasionally such operons contain several genes that are required to constitute one enzyme. To confirm that slr1737 is not required for phytyl prenyltransferase activity, phytyl prenyltransferase was measured in extracts  
5 from the *Synechocystis* slr1737 knockout mutant. Figure 29 shows that extracts from the *Synechocystis* slr1737 knockout mutant still contain phytyl prenyltransferase activity. The molecular organization of genes in *Synechocystis* 6803 is shown in A. Figures B and C show HPLC traces (normal phase HPLC) of reaction products obtained with membrane preparations from *Synechocystis* wild type and slr1737-  
10 membrane preparations, respectively.

The fact that slr1737 is not required for the PPT activity provides additional data that ATPT2 and slr1736 encode phytyl prenyltransferases.

#### 6C *Synechocystis* Knockouts

15 *Synechocystis* 6803 wild type and *Synechocystis* slr1737 knockout mutant were grown photoautotrophically. Cells from a 20 ml culture of the late logarithmic growth phase were harvested and extracted with ethanol. Extracts were separated by isocratic normal-phase HPLC using a Hexane/Methyl-t-butyl ether (95/5) and a Zorbax silica column, 4.6 x 250 mm. Tocopherols and tocopherol intermediates were  
20 detected by fluorescence (excitement 290 nm, emission 336 nm) (Figure 30).

Extracts of *Synechocystis* 6803 contained a clear signal of alpha-tocopherol. 2,3-Dimethyl-5-phytylplastoquinol was below the limit of detection in extracts from the *Synechocystis* wild type (C). In contrast, extracts from the *Synechocystis* slr1737 knockout mutant did not contain alpha-tocopherol, but contained 2,3-dimethyl-5-  
25 phytylplastoquinol (D), indicating that the interruption of slr1737 has resulted in a block of the 2,3-dimethyl-5-phytylplastoquinol cyclase reaction.

Chromatograms of standard compounds alpha, beta, gamma, delta-tocopherol and 2,3-dimethyl-5-phytylplastoquinol are shown in A and B. Chromatograms of extracts from *Synechocystis* wild type and the *Synechocystis* slr1737 knockout mutant



are shown in C and D, respectively. Abbreviations: 2,3-DMPQ, 2,3-dimethyl-5-phytylplastoquinol.

6D. Incubation with Lysozyme treated *Synechocystis*

5        *Synechocystis* 6803 wild type and slr1737 knockout mutant cells from the late logarithmic growth phase (approximately 1g wet cells per experiment in a total volume of 3 ml) were treated with Lysozyme and subsequently incubated with S-adenosylmethionine, and phytylpyrophosphate, plus radiolabelled homogentisic acid. After 17h incubation in the dark at room temperature the samples were extracted  
10 with 6 ml chloroform / methanol (1/2 v/v). Phase separation was obtained by the addition of 6 ml 0.9% NaCl solution. This procedure was repeated three times. Under these conditions 2,3-dimethyl-5-phytylplastoquinol is oxidized to form 2,3-dimethyl-5-phytylplastoquinone.

The extracts were analyzed by normal phase and reverse phase HPLC. Using  
15 extracts from wild type *Synechocystis* cells radiolabelled gamma-tocopherol and traces of radiolabelled 2,3-dimethyl-5-phytylplastoquinone were detected. When extracts from the slr1737 knockout mutant were analyzed, only radiolabelled 2,3-dimethyl-5-phytylplastoquinone was detectable. The amount of 2,3-dimethyl-5-phytylplastoquinone was significantly increased compared to wild type extracts. Heat  
20 treated samples of the wild type and the slr1737 knockout mutant did not produce radiolabelled 2,3-dimethyl-5-phytylplastoquinone, nor radiolabelled tocopherols. These results further support the role of the slr1737 expression product in the cyclization of 2,3-dimethyl-5-phytylplastoquinol.

25 6E. *Arabidopsis* Homologue to slr1737

An *Arabidopsis* homologue to slr1737 was identified from a BLASTALL search using *Synechocystis* sp 6803 gene slr1737 as the query, in both public and proprietary databases. SEQ ID NO:109 and SEQ ID NO:110 are the DNA and

translated amino acid sequences, respectively, of the *Arabidopsis* homologue to slr1737. The start is found at the ATG at base 56 in SEQ ID NO:109.

The sequences obtained for the homologue from the proprietary database differs from the public database (F4D11.30, BAC AL022537), in having a start site  
5 471 base pairs upstream of the start identified in the public sequence. A comparison of the public and proprietary sequences is provided in Figure 31. The correct start correlates within the public database sequence is at 12080, while the public sequence start is given as being at 11609.

Attempts to amplify a slr1737 homologue were unsuccessful using primers  
10 designed from the public database, while amplification of the gene was accomplished with primers obtained from SEQ ID NO:109.

Analysis of the protein sequence to identify transit peptide sequence predicted two potential cleavage sites, one between amino acids 48 and 49, and the other between amino acids 98 and 99.

15

#### 6F. slr1737 Protein Information

The slr1737 orf comprises 363 amino acid residues and has a predicted MW of 41kDa (SEQ ID NO: 39). Hydropathic analysis indicates the protein is hydrophilic (Figure 32).

20 The *Arabidopsis* homologue to slr1737 (SEQ ID xx) comprises 488 amino acid residues, has a predicted MW of 55kDa, and has a putative transit peptide sequence comprising the first 98 amino acids. The predicted MW of the mature form of the *Arabidopsis* homologue is 44kDa. The hydropathic plot for the *Arabidopsis* homologue also reveals that it is hydrophilic (Figure 33). Further blast analysis of  
25 the *Arabidopsis* homologue reveals limited sequence identity (25 % sequence identity) with the beta-subunit of respiratory nitrate reductase. Based on the sequence identity to nitrate reductase, it suggests the slr1737 orf is an enzyme that likely involves general acid catalysis mechanism.

Investigation of known enzymes involved in tocopherol metabolism indicated that the best candidate corresponding to the general acid mechanism is the tocopherol cyclase. There are many known examples of cyclases including, tocopherol cyclase, chalcone isomerase, lycopene cyclase, and aristolochene synthase. By further  
5 examination of the microscopic catalytic mechanism of phytoplastoquinol cyclization, as an example, chalcone isomerase has a catalytic mechanism most similar to tocopherol cyclase. (Figure 34).

Multiple sequence alignment was performed between slr1737, slr1737 *Arabidopsis* homologue and the *Arabidopsis* chalcone isomerase (Genbank:P41088)  
10 (Figure 35). 65% of the conserved residues among the three enzymes are strictly conserved within the known chalcone isomerases. The crystal structure of alfalfa chalcone isomerase has been solved (Jez, Joseph M., Bowman, Marianne E., Dixon, Richard A., and Noel, Joseph P. (2000) "Structure and mechanism of the evolutionarily unique plant enzyme chalcone isomerase". *Nature Structural Biology*  
15 7: 786-791.) It has been demonstrated tyrosine (Y) 106 of the alfalfa chalcone isomerase serves as the general acid during cyclization reaction (Genbank: P28012). The equivalent residue in slr1737 and the slr1737 *Arabidopsis* homolog is lysine (K), which is an excellent catalytic residue as general acid.

The information available from partial purification of tocopherol cyclase from  
20 *Chlorella protothecoides* (U.S. Patent No. 5,432,069), *i.e.*, described as being glycine rich, water soluble and with a predicted MW of 48-50kDa, is consistent with the protein informatics information obtained for the slr1737 and the *Arabidopsis* slr1737 homologue.

All publications and patent applications mentioned in this specification are  
25 indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claim.

## CLAIMS

What is claimed is:

1. An isolated nucleic acid sequence encoding a tocopherol cyclase.
- 5 2. An isolated nucleic acid sequence according to Claim 1, wherein said tocopherol cyclase is active in the cyclization of 2,3-dimethyl-5-phytylplastoquinol to tocopherol.
3. An isolated nucleic acid sequence according to Claim 1, wherein said tocopherol cyclase is active in the cyclization of 2,3-dimethyl-5-geranylgeranylplastoquinol to tocotrienol.
- 10 4. An isolated DNA sequence according to Claim 1, wherein said nucleic acid sequence is isolated from a eukaryotic cell source.
5. An isolated DNA sequence according to Claim 4, wherein said eukaryotic cell source is selected from the group consisting of mammalian, nematode, fungal, and plant cells.
6. The DNA encoding sequence of Claim 5 wherein said tocopherol cyclase protein is from  
15 *Arabidopsis*.
7. The DNA encoding sequence of Claim 6 wherein said tocopherol cyclase protein is encoded by a sequence of SEQ ID NO:109.
8. The DNA encoding sequence of Claim 7 wherein said tocopherol cyclase protein has an amino acid sequence of SEQ ID NO:110.
- 20 9. The DNA encoding sequence of Claim 4 wherein said tocopherol cyclase protein is from a source selected from the group consisting of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.
10. An isolated DNA sequence according to Claim 4, wherein said prokaryotic source is a *Synechocystis* sp.
- 25 11. The DNA encoding sequence of Claim 10 wherein said tocopherol cyclase protein is encoded by a sequence of SEQ ID NO:38.
12. The DNA encoding sequence of Claim 10 wherein said tocopherol cyclase protein has an amino acid sequence of SEQ ID NO:39.

13. A nucleic acid construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a nucleic acid sequence encoding a tocopherol cyclase, and a transcriptional termination region.
14. A nucleic acid construct according to Claim 13, wherein said nucleic acid sequence  
5 encoding tocopherol cyclase is obtained from an organism selected from the group consisting of a eukaryotic organism and a prokaryotic organism.
15. A nucleic acid construct according to Claim 14, wherein said nucleic acid sequence encoding tocopherol cyclase is obtained from a plant source.
16. A nucleic acid construct according to Claim 15, wherein said nucleic acid sequence  
10 encoding tocopherol cyclase is obtained from a source selected from the group consisting of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.
17. A nucleic acid construct according to Claim 13, wherein said nucleic acid sequence encoding tocopherol cyclase is obtained from a *Synechocystis* sp.
18. A plant cell comprising the construct of 13.
- 15 19. A plant comprising a cell of Claim 18.
- 20 A feed composition produced from a plant according to Claim 19.
21. A seed comprising a cell of Claim 18.
- 22 Oil obtained from a seed of Claim 21.
23. A natural tocopherol rich refined and deodorised oil which has been produced by  
20 a method of treating an oil according to Claim 22 by distilling under low pressure and high temperature, wherein said refined oil has reduced free fatty acids and a substantial percentage of tocopherol present in the pretreated oil.
24. A refined oil according to claim 23, wherein the pretreated oil is crude or pre-treated soybean oil.
- 25 25. A refined oil according to claim 23, wherein the refined oil is degummed and bleached.
26. A method for the alteration of the isoprenoid content in a host cell, said method comprising; transforming said host cell with a construct comprising as operably linked

components, a transcriptional initiation region functional in a host cell, a nucleic acid sequence encoding tocopherol cyclase, and a transcriptional termination region, wherein said isoprenoid compound selected from the group of tocopherols and tocotrienols .

27. The method according to Claim 26, wherein said host cell is selected from the group  
5 consisting of a prokaryotic cell and a eukaryotic cell.

28. The method according to Claim 27, wherein said prokaryotic cell is a *Synechocystis* sp.

29. The method according to Claim 27, wherein said eukaryotic cell is a plant cell.

30. The method according to Claim 29, wherein said plant cell is obtained from a plant  
selected from the group consisting of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, ,  
10 leek, cotton, and tomato.

31. A method for producing an isoprenoid compound of interest in a host cell, said method comprising obtaining a transformed host cell, said host cell having and expressing in its genome:

a construct having a DNA sequence encoding a tocopherol cyclase operably linked to a  
15 transcriptional initiation region functional in a host cell,  
wherein said isoprenoid compound selected from the group of tocopherols and tocotrienols.

32. The method according to Claim 31, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.

33. The method according to Claim 32, wherein said prokaryotic cell is a *Synechocystis* sp.


20 34. The method according to Claim 32, wherein said eukaryotic cell is a plant cell.

35. The method according to Claim 34, wherein said plant cell is obtained from a plant selected from the group consisting wherein said compound selected from the group of *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.

36. A method for increasing the biosynthetic flux in a host cell toward production of  
25 an isoprenoid compound, said method comprising; transforming said host cell with a construct comprising as operably linked components, a transcriptional initiation region functional in a host cell, a DNA encoding a tocopherol cyclase, and a transcriptional termination region, wherein said isoprenoid compound selected from the group of tocopherols and tocotrienols,.

37. The method according to Claim 36, wherein said host cell is selected from the group consisting of a prokaryotic cell and a eukaryotic cell.
38. The method according to Claim 37, wherein said prokaryotic cell is a *Synechocystis* sp.
39. The method according to Claim 37, wherein said eukaryotic cell is a plant cell.
- 5 40. The method according to Claim 39, wherein said plant cell is obtained from a plant selected from the group consisting *Arabidopsis*, soybean, corn, rice, wheat, leek canola, , leek, cotton, and tomato.
41. The method according to Claim 39, wherein said transcriptional initiation region is a seed-specific promoter.





## Figure 1

2/40





Figure 2



### Figure 3

4/40




Figure 4

5/40




Figure 5

6/40




Figure 6

7/40




Figure 7

8/40




Figure 8



9/40




Figure 9

10/40




Figure 10

11/40




Figure 11

12/40




Figure 12

13/40




Figure 13

14/40




Figure 14

15/40




Figure 15

16/40




Figure 16



17/40




Figure 17

18/40




Figure 18

19/40




Figure 19

20/40




Figure 20



22/40

ATPT2	:	20	*	40	*	60	*	80
SLR1736	:	MESLLSSSLVSAAGGFCWKQNLKLSLSEIRVLRCDSKVKVAKPKFRNNLVRDQGGSSLLLYPKHSRFRVNATAGQ						
ATPT3	:	MAFFGLSRVSRLLKSSVSTPSSSALLQSHKSLSNPVTHYTNPTKCYPSWNDNYQVWSKGRELHQKFFGVGWNRYRLICGMSSS						
SLR0926	:	MWERSVYRFSSRISVSSSLPNRLPWSRELCAVNSFSQP						
ATPT4	:	PVSSTESTAKLIGITGVRSDANRVFATA						
SLR1899	:	MTSILNTVSTIHSRVTSDRVGLSLRNSDSVEFT						
ATPT12	:	RRRSSTLIYESPGRFVVRAAETDT						
SLR0056	:	MSDT						
ATPT8	:	MVLA						
SLR1518	:	MTES						

ATPT2	:	100	*	120	*	140	*	160	*
SLR1736	:	PEAFDSNSKQK							
ATPT3	:	SSVLEKPKKDDKEKSDGVVVKASWDLXLPPEVRGYAKLARLDKPIGTWELAWPCWMS							
SLR0926	:	MVAQTPSSP							
ATPT4	:	TAAATATATTG							
SLR1899	:	TKIHRQDSMG							
ATPT12	:	DKVKSQTPDKAP							
SLR0056	:	QNT-GQNOAKA							
ATPT8	:	EVPKLASAERY							
SLR1518	:	SPLAPSTAPAT							

ATPT2	:	200	*	220	*	240	*	260
SLR1736	:	AVVAALMNVYIVG						
ATPT3	:	GALL						
SLR0926	:	GTEA						
ATPT4	:	TMTI						
SLR1899	:	GTEA						
ATPT12	:	MTSGPC						
SLR0056	:	MLSGPL						
ATPT8	:	GIAE						
SLR1518	:	SATA						

Figure 22A

23/40

ATPT2	280	*	300	*	320	*	340	*	360	*	380	*	400	*	420	*	440	*	460	*	480	*
SLR1736	WKR-FALVAAMCILAVRAIIQIAFYH-IQTHVFRPIETFRPPIFATAFMSFSSVVIAMFKDIPDIKGGKI-FEIRSFSTLIG-QKR	313																				
ATPT3	LKR-PSLLAALCILTVRGIVNGLFIF-FRIGLGYPTLITPFW-ETLFIILVAVAIAMFKDVPDMGGRQ-FKIQITLQIG-KQN	218																				
SLR0926	SXP-LMKRFTWPQAFGLTNWGATIG-WT-RVKGSAFSPLEFYLGVCKDQVYDIYAHQDKDQVK-FCGKSTAFRFQ-DNT	328																				
ATPT4	AMP-GAKRVFPVQOLLSIANGFAVHIS-SS-PAIADDAITW-ANGATVMTLGFDTVYAMADRDERR-FCUNSSALFFQ-QYV	213																				
SLR1899	VMT-PLXQLHPINTWGAUVGAIPPIIG-WA-WASQGLSYNSMIPPAALFYQHPHFMAHALHLCRNDYAA-CGYQTSFDFP--S	294																				
ATPT12	VATHMLKRHTAQNIVIGGAAGSIPPIIG-WA-WATBDSWTPW-EPALIFLAMPHPHFMAHALMIKODDYAQ-NNWPMPPVIAGEEKT	220																				
SLR0056	INS-APPLKQNGWGNFALGASYHISLPWAGQAFPTIPDVW-ETLLYSIAGGIAVNDKSVGGRQ-FCGOSPVAFQ-TET	308																				
ATPT8	IS-APPLKQNGWGNFALGASYHISLPWAGQAFPTIPDVW-ETLLYSIAGGIAVNDKSVGGRQ-FCGKSEPMWFG-IGT	242																				
SLR1518	EITSSTEQRYSNDYYOKTYTKTASLSSNSCKAVALLTQGAELVAMLAFFEYGRNLGAFQIIDDILDTGTSASCKGSLSDIRH--GV	231																				
	TGQPPFRLGYLGLGELICITTFGLAI-AAAYYSQSQSFSWNIT-PSVFVGISFALILFCSHFHQVDSLA-ACKKSPIVRIQ-TKL	223																				
ATPT2	360	*	380	*	400	*	420	*	440	*	460	*	480	*	500	*	520	*	540	*	560	*
SLR1736	VFWTQITLQWYARAIEVGGATSPFIWSKVISVGHVITATTWARAKSVDSLSSKTEHTS--CYNMIWKFFYAEYVILPFLK----	393																				
ATPT3	VFRGTHILATGCVLAMAATWGAAMPLNTAFPTVSHLCILALFWRSRDVHLESKTEHTAS--FYQIWKFFLEYILYPLAWLPNFS	304																				
SLR0926	KLMTGFGTASIGFTALSGFSADLGWQYVASAAASGQGWQIGTADLSSGADCS-----RQVSNKWFQGFSPGVNQRSFQ	407																				
ATPT4	GEANGFFALTIGCFYTGNDMLNPLYWLSAQAQ--GQWFOYIQLSAPTPEP-KQY-----GOIQGONIIIGFVILAGMIGWL--	292																				
SLR1899	GKRLAAVARNCFYNIPIGFGAYDWGLTSSWFCEBETLTLAAATAFSRYDRTWKKA-----RKNVHASLFLPWFNSGMLHRVSND	379																				
ATPT12	VSOHWYSLLVVPFSLIVVPLHQGLILYLAHAIIT--GGQFLVKAWLKQAPGDRDIA-----RGLKFSFYIMLICTAMVDDSLPVT	303																				
SLR0056	AKWTCGAIIDITQLSVAGYLAASKPYVALMAVALA--IPQVFPQFKYFLKDPVKYDK-----YQASAQPFVLVLCITATASQH----	387																				
ATPT8	AANGCQIMIDVFQAGIAGYLTVHQQLYATITLIL--PQTFQDMYFLNPLENDYK-----YQASAQPFVLVLCITATASQH----	324																				
SLR1518	ITAPITFAMEEFPQUREVDOVEKDPNRVDIAEXTGKSKGTORARELAHEHANLAAAAGISLPETDNEDVKKSRRAIIDLTHRVITRN	320																				
	GSOVLTLSVSLVLTATAGVTCQAPWQITLITASAPWAVQVIRHVGVQHDQPEQVSNCK--FLAVNLHFFSGMLAAGYGNWAGLG--	307																				
ATPT2	-----																					
SLR1736	NTIF-----																					
ATPT3	-----																					
SLR0926	-----																					
ATPT4	NOQLVEEAGLTNSVSGEVTKQRKKRVAQPPVAYASAPFFFLPAPFISP	431																				
SLR1899	--HQLVAQMGTLLIG-----	316																				
ATPT12	-----																					
SLR0056	-----																					
ATPT8	K-----	321																				
SLR1518	-----																					

Figure 22B

24/40




Figure 23



25/40




Figure 24

26/40




Figure 25




Figure 26

28/40





Figure 27

29/40

### Total tocopherol in Napin ATPT2 Canola Seed

**Figure 28**

30/40




Figure 29

31/40




Figure 30

32/40

### Alignment 1

[illegible]

**Figure 31A**



33/40

```
Query-      11595 CAGAGAGTTTTTTTATGTTGATAACTTATTGTTTAACTTTGAAAAATGCAGATTA
ATCEA4C371+  299 -----ATA
PIR:T04448    6 R E F F F L W L I T Y C L T F E K C R Y

Query-      11535 CCATTCGATGGAACACCTCGGAAGTTCTTCGAGGGATGGTATTTTCAGGGTTTCCATCCC
ATCEA4C371+  302 CCATTCGATGGAACACCTCGGAAGTTCTTCGAGGGATGGTATTTTCATCCATCCC
PIR:T04448    26 H F D G T P R K F F E G W Y F S I P

Query-      11475 AGAGAAGAGGGAGAGTTTTTGTGTTTATGTATTCTGTGGAGAATCCTGCATTCGGCAGAG
ATCEA4C371+  362 AGAGAAGAGGGAGAGTTTTTGTGTTTATGTATTCTGTGGAGAATCCTGCATTCGGCAGAG
PIR:T04448    46 E K R E S F C F M Y S V E N P A F R Q S

Query-      11415 TTTGTCACCATTTGGAAGTGGCTCTATATGGACCTAGATTCACTGGTGTGGAGCTCAGAT
ATCEA4C371+  422 TTTGTCACCATTTGGAAGTGGCTCTATATGGACCTAGATTCACTGGTGTGGAGCTCAGAT
PIR:T04448    66 L S P L E V A L Y G P R F T G V G A Q I

Query-      11355 TCTTGGCGCTAATGATAAATATTTATGCCAATACGAACAAGACTCTCACAATTTCTGGGG
ATCEA4C371+  482 TCTTGGCGCTAATGATAAATATTTATGCCAATACGAACAAGACTCTCACAATTTCT
PIR:T04448    86 L G A N D K Y L C Q Y E Q D S H N F W G
ATCEA4C371+  Exon      11538      11301 Confidence: 100 100

Query-      11295 AGGTAACCTCCTTGACCCTTAAATGCTGTGTCATGACAATAAGAAATCATATCTGAGTCT
ATCEA4C371+  537 -----
PIR:T04448    106 D
PIR:T04448    Exon      11609      11294 Confidence: 100 100

Query-      11235 TTTCTCTACTTCTAGTACTAATGTTTCGTTATTGTTGTTAAAGATCTAAGTCTTATCTGAA
PIR:T04448    107 -----

Query-      11175 TTTTGTACATTTTGGTTCTGCTGCTTTCTCAACATGAATTTGTATATATGACTTTAAAG
PIR:T04448    107 -----

Query-      11115 ATTGCTTACCTAAAGTTTTTACTCATGCATAGATCGACATGAGCTAGTTTGGGGAATAC
PIR:T04448    107 -----R H E L V L G N T

Query-      11055 TTTTAGTGCTGTGCCAGGCGCAAAGGCTCCAAACAAGGAGTTCCACCAGAGGTTCTCAC
PIR:T04448    116 F S A V P G A K A P N K E V P P E
PIR:T04448    Exon      11083      11004 Confidence: 96 100
```

Figure 31B

34/40

Query- 10995 TCCTCCCTTGGTTACTTTGTTATCTGTTAAATAGTTTCCAATTGTATCCGGATAGT  
PIR:T04448 133

Query- 10935 GTTCTACTTCTCCTTGTAGAAAATCTCAAGTTTTGTTACTCTGCTATTCTCTTGGATG  
PIR:T04448 133

Query- 10875 TTGATTTGTAAAGCATGTCGTTTTATTGTAGGAATTTAACAGAAGAGTGTCCGAAGGGTT  
PIR:T04448 133  
E F N R R V S E G F

Query- 10815 CCAAGCTACTCCATTTGGCATCAAGGTCACATTGCGATGATGGCCGGTAATTATATGA  
PIR:T04448 143  
Q A T P F W H Q G H I C D D G R  
PIR:T04448 Exon 10844 10768 Confidence: 100 100

Query- 10755 TTCTATGCACAACAAGAATTCACTATATTATAAATATTGGATATTGAGTATTTTGTGA  
PIR:T04448 159

Query- 10695 AAATTCTGTGTTTAAATCTGACTTGACTTGTGTTGTCAGTACTGACTATGCCGAAACTG  
PIR:T04448 159  
T D Y A E T V

Query- 10635 TGAAATCTGCTCGTTGGAGTATAGTACTCGTCCCGTTACGGTTGGGGTGATGTTGGGG  
PIR:T04448 166  
K S A R W E Y S T R P V Y G W G D V G A

Query- 10575 CCAAACAGAGTCAACTGCAGGCTGGCCTGCAGCTTTTCTCTGTATTGAGCCTCATTTGGC  
PIR:T04448 186  
K Q K S T A G W P A A F P V F E P H W Q

Query- 10515 AGATATGCATGGCAGGAGGCCCTTCCACAGGTGTGAGCTTTGCTTGATTGACTTAAAGTT  
PIR:T04448 206  
I C M A G G L S T G  
PIR:T04448 Exon 10655 10486 Confidence: 96 100

Query- 10455 AATAAATAGACGGTTAAGTTTACTTGCCTAGTACTAACAGAAAATTAAGAAAGAAACCAC  
PIR:T04448 216

Query- 10395 CCTCTTCTATCAGCAGAACTGCTATTGTAGTTCTTATTTTTCTCTTGTATTGCAGG  
PIR:T04448 216

Query- 10335 GTGGATAGAATGGGCGGTGAAAGGTTTGTAGTTTCGGGATGCACCTTCTTATTCAGAGAA  
PIR:T04448 216  
W I E W G G E R F E F R D A P S Y S E K

Query- 10275 GAATTGGGGTGGAGGCTTCCCAAGAAAATGGTTTGGGGTAAACATTTTCATCCTTTTGCT  
PIR:T04448 236  
N W G G G F P R K W F W  
PIR:T04448 Exon 10336 10239 Confidence: 96 100

Figure 31C

35/40

```
Query- 10215 ACATTCTTGTTCAGACTTTAGTTAGCTAGTGGACCTGTGTATACACCCACATGTAGTA
PIR:T04448 248 -----

Query- 10155 TACTTGTGTGATAGCTTTATTTGTCAATGTCTCTTTACAGGTCCAGTGTAAATGTCTTTGA
PIR:T04448 248 -----
V Q C N V F E

Query- 10095 AGGGGCAACTGGAGAAGTTGCTTTAACCAGGCGGGTTGAGGCAATTGCCTGGATT
PIR:T04448 255 G A T G E V A L T A G G G L R Q L P G L

Query- 10035 GACTGAGACCTATGAAAATGCTGCACTGGTATGCACTTATAAGATCTTCTTAAGCAATGA
PIR:T04448 275 T E T Y E N A A L
PIR:T04448 Exon 10115 10008 Confidence: 100 100

Query- 9975 CAGTGAGTATTAGAAGGCAGATAGTTTACAAAAGCTCTGGGCCCTTGTAATCTGCAGGT
PIR:T04448 284 -----
V

Query- 9915 TTGTGTACACTATGATGGAAAAATGTACGAGTTTGTTCCTTGGAAATGGTGTGTAGATG
PIR:T04448 285 C V H Y D G K M Y E F V P W N G V V R W
GSDB:S:495- 532 -----
tagatg

Query- 9855 GGAAATGTCTCCCTGGGG TTATTGGTATATAACTGCAGAGAACGAAAACCATGTGGTAA
PIR:T04448 305 E M S P W G Y W Y I T A E N E N H V
GSDB:S:495- 526 ggaaat tctccctgggggttattggtatataactgcagagancgNaaacctgtg
PIR:T04448 Exon 9917 9801 Confidence: 100 100
GSDB:S:495- Exon 9861 9801 Confidence: 93 93

Query- 9796 ATTTGTTTACTAGTTTCATTCACTTTACTTTTGACATCATATCATTCCCTTATGGCTA
PIR:T04448 323 -----
GSDB:S:495- 471 -----

Query- 9736 GATTCCAACACCCGATGAATGCTTGTGACAGGTGGAAGTAGAGGCAAGAACAAATGAAG
PIR:T04448 323 -----
V E L E A R T N E A
GSDB:S:495- 471 -----
gtggaactagaggcNagaacaaatgaag

Query- 9676 CGGGTACACCTCTGCGTGCTCTACCACAGAAGTTGGGCTAGCTACGGCTTGCAGAGATA
PIR:T04448 333 G T P L R A P T T E V G L A T A C R D S
GSDB:S:495- 443 cgggtacacctctgcgtgctcctaccacagaagttgggctagctacggcttgcagagata

Query- 9616 GTTGTTACGGTGAATTGAAGTTGCAGATATGGGAACGGCTATATGATGGAAGTAAAGGCA
PIR:T04448 353 C Y G E L K L Q I W E R L Y D G S K G K
GSDB:S:495- 383 gttgttacggtgaattgaagttgcagatatgggaacggctatatgatggaagtaaaggca
```

Figure 31D

36/40

Query- 9556 AGGTATGTATGCTAATGTGATCCAATCCCTGTAGTTAAAAGTCTTAACAAATCCTAAGGC  
PIR:T04448 373 ::  
GSDB:S:495- 323 ||----- L K V L T N P K A  
PIR:T04448 323 ag  
GSDB:S:495- Exon 9704 9555 Confidence: 100 100  
Exon 9704 9555 Confidence: 98 100

Query- 9496 AGTGAAGAAGATTATGAACGTTTGTATTGGTTAACAATGATGCAGGTGATATTAGAGAC  
PIR:T04448 382 ::  
GSDB:S:495- 321 V K E D Y E R L L W L T M M Q V I L E T  
gtgatattagagac

Query- 9436 AAAGAGCTCAATGGCAGCAGTGGAGATAGGAGGAGGACCGTGGTTGGGACATGGAAAGG  
PIR:T04448 402 ::  
GSDB:S:495- 307 K S S M A A V E I G G G P W F G T W K G  
aaagagctcaatggcancagtgagataggaggaggaccgtggtttgggacatggaaagg

Query- 9376 AGATACGAGCAACACGCCCGAGCTACTAAAACAGGCTCTTCAGGTCCCATTGGATCTTGA  
PIR:T04448 422 ::  
GSDB:S:495- 247 D T S N T P E L L K Q A L Q V P L D L E  
agatacgagcaacacgcccagctactaaacaggctcttcaggtcccattggatcttga

Query- 9316 AAGCGCCTTAGGTTTGGTCCCTTCTTCAAGCCACCGGGTCTG TAA  
(stop) 442 ::  
PIR:T04448 187 S A L G L V P F F K P P G L  
GSDB:S:495- aagcgcccttaggtttgtcccttcttcaagccacgggtctgtaacattgatgagtgt  
PIR:T04448 Exon 9522 9274 Confidence: 100 100

Query- 9256 [REDACTED]  
PIR:T04448 456 [REDACTED]  
GSDB:S:495- 127 ttgtttgttgatagagaccatgtgatgaatgaagccttagtcatgtcattgctagcttc

Query- 9196 ACTATTATGTATGTATGATTTTAGTTCGTTTCGGTCTTGTGGTAAATGATACGGGCCAGT  
GSDB:S:495- 67 actattatgtatgtatgatttagttcgttcggtccttgtggtaaatgatacgggccagt

Query- 9136 GTAAAGTCTAGTTCAATAAAGCCTTGAGTCGCATAATTCAATTTCAAATTGCATC  
GSDB:S:495- 7 gtaaagt  
GSDB:S:495- Exon 9450 9130 Confidence: 98 100

ATCEA4C37145\_1 3063693/emb|CAA18584.1| 4.0e-43 (AL022537) putative protein  
[Arabidopsis thaliana]  
PIR:T04448 sPIR-T04448 shypothetical protein F4D11.30 - Arabidopsis thaliana;  
g3063693|emb|CAA18584.1 (AL022537) putative protein [Arabidopsis thaliana]\_F4D11.30  
GSDB:S:4955486|AI995392|AI995392|701673779 A. thaliana, Columbia Col-0, inflorescence-  
1 Arabidopsis thaliana cDNA clone 701673779, mRNA sequence.

Figure 31E

37/40




Figure 32

38/40




Figure 33

39/40




Figure 34

40/40

slr1737_SYNSP_S74814	-----M
slr1737_ARATH_T04448	MEIRSLIVSMNPNLSSFELSRPVSPLTRSLVPFRSTKLVPRSISRVSASI
CFI_ARATH_P41088	-----
slr1737_SYNSP_S74814	KFP-----PHSGYHWQGS-PFFEGWYVRL
slr1737_ARATH_T04448	STPNSETDKISVKPVYVPTSPNRELRTPHSGYHFDGTPRKFFEGWYFRVS
CFI_ARATH_P41088	-----
slr1737_SYNSP_S74814	LPOSGESFAFMYSIENPASDHHYGGGAVQILGPATK-----KOENQEDQLV
slr1737_ARATH_T04448	IPEKRESFCFMYSVENPAFRQSLSPLEVALYGPFRFTGVGAQILGANDKYL
CFI_ARATH_P41088	MSSSNACASPSPPFA-----VTKLHVDSV-
slr1737_SYNSP_S74814	WRTFPSVKKFWASPRQFALG-HWGKCRDNRO-AKPLLSEEFFATVKEGYQ
slr1737_ARATH_T04448	CQYEQDSHNEWGDRLVLGNTFSVAVPGAKAPNKEVPPEEFNRRVSEGFQ
CFI_ARATH_P41088	--TFVPSVKSPASSNPLFLG-GAGVRGLDIQ-GK-----FVIFTVIGVY
slr1737_SYNSP_S74814	IHQNHQHQQIIHGDR-----HCRWQFTVEPEVTWGSNRFPRATAGW
slr1737_ARATH_T04448	ATPFWHQGHICDDGRDYAETVKSARWEYSTRPVYWGVDVGAKQKSTAGW
CFI_ARATH_P41088	LEGNVPSLSV-----KWKGKTTEELTESIPFFREIVTGAF
slr1737_SYNSP_S74814	LSFLPLFDPGWQILLAQGRAHGWLKWQREQYEFDHALVYAEKNWGHSEFPS
slr1737_ARATH_T04448	PAAFPVFEPHWQICMAGGLSTGWIEWGGERFERDAPSYSEKNWGGGFPR
CFI_ARATH_P41088	EKFIKVT-----M-----KLPLTGQQYSEKVTENC
slr1737_SYNSP_S74814	RWFWLQANYFPDHPG-LSVTAAGGERIVLGRPE---EVALIGLHHQGNFY
slr1737_ARATH_T04448	KWFWVQCNVFEAGATGEVALTAGGGLRQLPGLTETYENALVCVHYDGKMY
CFI_ARATH_P41088	VAIWKQLGLTYDCEA-KAV-----EKFLEIFKE---ET-----
slr1737_SYNSP_S74814	EFGPGHGTVTWQVAPWGRWQLKASNDRYWVKLSGKTDKKGSLVHTP-TAQ
slr1737_ARATH_T04448	EFVPWNGVVRWEMSPWGYWYITAENENHVVELEARTNEAGTPLRAPTTEV
CFI_ARATH_P41088	-FPPG-SSILFALSPTGSLTVAFSKDDS-IPETGIAVIENKLLAEA-VLE
slr1737_SYNSP_S74814	GLQLNCRDTRGYLYLQLGSGVGHG-----LIVQGETDTAGLEVGG-----
slr1737_ARATH_T04448	GLATACRDCYSELKQLQIWERLYDGSKGKVILETKSSMAAVEIGGGPWFG
CFI_ARATH_P41088	--SIIGKNGVSPGTRLSVAERLSQ-----LMMKNKDEKEVSDHSL-----
slr1737_SYNSP_S74814	-----DWGLTEENLSKKT-----VPF-----
slr1737_ARATH_T04448	TWKGDTSNTPELLKQALQVPLDLESALGLVPFFKPPGL
CFI_ARATH_P41088	-----EEKLAKEN-----

Figure 35



## SEQUENCE LISTING

<110> Subramaniam, Sai  
 5 Slater, Steven  
 Karberg, Katherine  
 Chen, Ridong  
 Valentin, Henry  
 Huang Wong, Yun-Hua  
 10  
 <120> Nucleic Acid Sequences Involved in  
 Tocopherol Synthesis  
  
 <130> MOCO.008.00WO  
 15  
 <150> 09/549,848  
 <151> 2000-04-15  
  
 <150> 09/688,069  
 20 <151> 2000-10-15  
  
 <160> 94  
  
 <170> FastSEQ for Windows Version 4.0  
 25  
 <210> 1  
 <211> 1182  
 <212> DNA  
 <213> Arabidopsis sp  
 30  
 <400> 1  
 atggagtctc tgctctctag ttcttctctt gtttccgctg ctggtgggtt ttgttggaag 60  
 aagcagaatc taaagctoca ctctttatca gaaatccgag ttctgcgttg tgattcgagt 120  
 aaagtgtgctg caaaaccgaa gtttaggaac aatcttgta ggcctgatgg tcaaggatct 180  
 35 tcattgttgt tgtatccaaa acataagtcg agatttcggg ttaatgccac tgcgggtcag 240  
 cctgaggctt tcgactcgaa tagcaaacag aagtctttta gagactcggt agatgcgttt 300  
 tacagggttt ctaggcctca tacagttatt ggcacagtgc ttagcatttt atctgtatct 360  
 ttcttagcag tagagaaggt ttctgatata tctcctttac ttttcactgg catcttgag 420

	gctgtgttg cagctctcat gatgaacatt tacatagttg ggctaaatca gttgtctgat	480
	gttgaaatag ataaggtaa caagccctat cttccattgg catcaggaga atattctggt	540
	aacaccggca ttgcaatagt agcttccttc tccatcatga gtttctggct tgggtggatt	600
	gttggttcat ggccattgtt ctgggctctt tttgtgagtt tcatgctcgg .tactgcatac	660
5	tctatcaatt tgccactttt acggtggaaa agatttgcatt tgggtgcagc aatgtgtatc	720
	ctcgctgtcc gagctattat tgttcaaato gccttttatac tacatattca gacacatgtg	780
	tttggagac caatcttggt cactaggcct cttattttcg ccactgcgtt tatgagcttt	840
	ttctctgtcg ttattgcatt gtttaaggat atacctgata tcgaaggga taagatatcc	900
	ggaatccgat cattctctgt aactctgggt cagaaacggg tgttttgac atgtgttaca	960
10	ctacttcaaa tggttacgc tgttgcaatt ctagttggag ccacatctcc attcatatgg	1020
	agcaaagtca tctcggttgt gggcatggt atactcgaa caactttgtg ggctcgagct	1080
	aagtcggttg atctgagtag caaaaccgaa ataacttcat gttatatgtt catatggaag	1140
	ctcttttatg cagagtactt gctgttacct ttttgaagt ga	1182
15	<210> 2	
	<211> 393	
	<212> PRT	
	<213> Arabidopsis sp	
20	<400> 2	
	Met Glu Ser Leu Leu Ser Ser Ser Ser Leu Val Ser Ala Ala Gly Gly	
	1 5 10 15	
	Phe Cys Trp Lys Lys Gln Asn Leu Lys Leu His Ser Leu Ser Glu Ile	
	20 25 30	
25	Arg Val Leu Arg Cys Asp Ser Ser Lys Val Val Ala Lys Pro Lys Phe	
	35 40 45	
	Arg Asn Asn Leu Val Arg Pro Asp Gly Gln Gly Ser Ser Leu Leu Leu	
	50 55 60	
	Tyr Pro Lys His Lys Ser Arg Phe Arg Val Asn Ala Thr Ala Gly Gln	
30	65 70 75 80	
	Pro Glu Ala Phe Asp Ser Asn Ser Lys Gln Lys Ser Phe Arg Asp Ser	
	85 90 95	
	Leu Asp Ala Phe Tyr Arg Phe Ser Arg Pro His Thr Val Ile Gly Thr	
	100 105 110	
35	Val Leu Ser Ile Leu Ser Val Ser Phe Leu Ala Val Glu Lys Val Ser	
	115 120 125	
	Asp Ile Ser Pro Leu Leu Phe Thr Gly Ile Leu Glu Ala Val Val Ala	
	130 135 140	
	Ala Leu Met Met Asn Ile Tyr Ile Val Gly Leu Asn Gln Leu Ser Asp	

145                      150                      155                      160  
 Val Glu Ile Asp Lys Val Asn Lys Pro Tyr Leu Pro Leu Ala Ser Gly  
                          165                      170                      175  
 Glu Tyr Ser Val Asn Thr Gly Ile Ala Ile Val Ala Ser Phe Ser Ile  
 5                      180                      185                      190  
 Met Ser Phe Trp Leu Gly Trp Ile Val Gly Ser Trp Pro Leu Phe Trp  
                          195                      200                      205  
 Ala Leu Phe Val Ser Phe Met Leu Gly Thr Ala Tyr Ser Ile Asn Leu  
                          210                      215                      220  
 10 Pro Leu Leu Arg Trp Lys Arg Phe Ala Leu Val Ala Ala Met Cys Ile  
                          225                      230                      235                      240  
 Leu Ala Val Arg Ala Ile Ile Val Gln Ile Ala Phe Tyr Leu His Ile  
                          245                      250                      255  
 Gln Thr His Val Phe Gly Arg Pro Ile Leu Phe Thr Arg Pro Leu Ile  
 15                      260                      265                      270  
 Phe Ala Thr Ala Phe Met Ser Phe Phe Ser Val Val Ile Ala Leu Phe  
                          275                      280                      285  
 Lys Asp Ile Pro Asp Ile Glu Gly Asp Lys Ile Phe Gly Ile Arg Ser  
                          290                      295                      300  
 20 Phe Ser Val Thr Leu Gly Gln Lys Arg Val Phe Trp Thr Cys Val Thr  
                          305                      310                      315                      320  
 Leu Leu Gln Met Ala Tyr Ala Val Ala Ile Leu Val Gly Ala Thr Ser  
                          325                      330                      335  
 Pro Phe Ile Trp Ser Lys Val Ile Ser Val Val Gly His Val Ile Leu  
 25                      340                      345                      350  
 Ala Thr Thr Leu Trp Ala Arg Ala Lys Ser Val Asp Leu Ser Ser Lys  
                          355                      360                      365  
 Thr Glu Ile Thr Ser Cys Tyr Met Phe Ile Trp Lys Leu Phe Tyr Ala  
                          370                      375                      380  
 30 Glu Tyr Leu Leu Leu Pro Phe Leu Lys  
                          385                      390

&lt;210&gt; 3

&lt;211&gt; 1224

35 &lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 3

atggcggtttt ttgggctctc ccgtgtttca agacgggttg tgaaatcttc cgtctccgta

60

actccatctt cttcctctgc tcttttgcaa tcacaacata aatccttgtc caatcctgtg 120  
 actaccatt acacaaatcc tttcactaag tgttatcctt catggaatga taattaccaa 180  
 gtagggagta aaggaagaga attgcatcag gagaagtttt ttggtggttg ttggaattac 240  
 agattaattt gtggaatgtc gtcgtcttct tcggttttgg agggaaagcc gaagaaagat 300  
 5 gataaggaga agagtgatgg tgttgttgtt aagaaagctt cttggataga tttgtattta 360  
 ccagaagaag ttagagggtta tgctaagctt gctcgattgg ataaacccat tggaacttgg 420  
 ttgcttgctg ggccttgat gtggtcgatt gcgttggtg ctgatectgg aagccttcca 480  
 agttttaaat atatggcttt atttggttgc ggagcattac ttottagagg tgctggttgt 540  
 actataaatg atctgcttga tcaggacata gatacaaagg ttgatectac aaaactaaga 600  
 10 cctatcgcca gtggtctttt gacaccattt caagggattg gatttctcgg gctgcagttg 660  
 cttttaggct tagggattct tctccaactt aacaattaca gccgtgtttt aggggcttca 720  
 tctttgttac ttgtcttttc ctaccactt atgaagaggt ttacattttg gcctcaagcc 780  
 tttttagggt tgaccataaa ctggggagca ttgttaggat ggactgcagt taaaggaagc 840  
 atagcaccat ctattgtact cctctctat ctctccggag tctgctggac ccttgtttat 900  
 15 gatactattt atgcacatca ggacaaagaa gatgatgtaa aagttggtgt taagtcaaca 960  
 gcccttagat tcggtgataa tacaagctt tggtaactg gatttggcac agcatccata 1020  
 ggttttcttg cactttcttg attcagtga gatctcgggt ggcaatatta cgcactcgtg 1080  
 gccgctgcat caggacagtt aggatggcaa ataggacag ctgacttata atctggtgct 1140  
 gactgcagta gaaaatttgt gtcgaacaag tggtttggtg ctattatatt tagtggagtt 1200  
 20 gtacttgga gaagttttca ataa 1224

&lt;210&gt; 4

&lt;211&gt; 407

&lt;212&gt; PRT

25 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 4

Met Ala Phe Phe Gly Leu Ser Arg Val Ser Arg Arg Leu Leu Lys Ser  
 1 5 10 15  
 30 Ser Val Ser Val Thr Pro Ser Ser Ser Ser Ala Leu Leu Gln Ser Gln  
 20 25 30  
 His Lys Ser Leu Ser Asn Pro Val Thr Thr His Tyr Thr Asn Pro Phe  
 35 40 45  
 Thr Lys Cys Tyr Pro Ser Trp Asn Asp Asn Tyr Gln Val Trp Ser Lys  
 35 50 55 60  
 Gly Arg Glu Leu His Gln Glu Lys Phe Phe Gly Val Gly Trp Asn Tyr  
 65 70 75 80  
 Arg Leu Ile Cys Gly Met Ser Ser Ser Ser Ser Val Leu Glu Gly Lys  
 85 90 95

Pro Lys Lys Asp Asp Lys Glu Lys Ser Asp Gly Val Val Val Lys Lys  
 100 105 110  
 Ala Ser Trp Ile Asp Leu Tyr Leu Pro Glu Glu Val Arg Gly Tyr Ala  
 115 120 125  
 5 Lys Leu Ala Arg Leu Asp Lys Pro Ile Gly Thr Trp Leu Leu Ala Trp  
 130 135 140  
 Pro Cys Met Trp Ser Ile Ala Leu Ala Ala Asp Pro Gly Ser Leu Pro  
 145 150 155 160  
 Ser Phe Lys Tyr Met Ala Leu Phe Gly Cys Gly Ala Leu Leu Leu Arg  
 10 165 170 175  
 Gly Ala Gly Cys Thr Ile Asn Asp Leu Leu Asp Gln Asp Ile Asp Thr  
 180 185 190  
 Lys Val Asp Arg Thr Lys Leu Arg Pro Ile Ala Ser Gly Leu Leu Thr  
 195 200 205  
 15 Pro Phe Gln Gly Ile Gly Phe Leu Gly Leu Gln Leu Leu Leu Gly Leu  
 210 215 220  
 Gly Ile Leu Leu Gln Leu Asn Asn Tyr Ser Arg Val Leu Gly Ala Ser  
 225 230 235 240  
 Ser Leu Leu Leu Val Phe Ser Tyr Pro Leu Met Lys Arg Phe Thr Phe  
 20 245 250 255  
 Trp Pro Gln Ala Phe Leu Gly Leu Thr Ile Asn Trp Gly Ala Leu Leu  
 260 265 270  
 Gly Trp Thr Ala Val Lys Gly Ser Ile Ala Pro Ser Ile Val Leu Pro  
 275 280 285  
 25 Leu Tyr Leu Ser Gly Val Cys Trp Thr Leu Val Tyr Asp Thr Ile Tyr  
 290 295 300  
 Ala His Gln Asp Lys Glu Asp Asp Val Lys Val Gly Val Lys Ser Thr  
 305 310 315 320  
 Ala Leu Arg Phe Gly Asp Asn Thr Lys Leu Trp Leu Thr Gly Phe Gly  
 30 325 330 335  
 Thr Ala Ser Ile Gly Phe Leu Ala Leu Ser Gly Phe Ser Ala Asp Leu  
 340 345 350  
 Gly Trp Gln Tyr Tyr Ala Ser Leu Ala Ala Ala Ser Gly Gln Leu Gly  
 355 360 365  
 35 Trp Gln Ile Gly Thr Ala Asp Leu Ser Ser Gly Ala Asp Cys Ser Arg  
 370 375 380  
 Lys Phe Val Ser Asn Lys Trp Phe Gly Ala Ile Ile Phe Ser Gly Val  
 385 390 395 400  
 Val Leu Gly Arg Ser Phe Gln

405

&lt;210&gt; 5

&lt;211&gt; 1296

5 &lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

&lt;400&gt; 5

	atgtggcgaa gatctgttgt ttctcgttta tcttcaagaa tctctgtttc ttcttcgtta	60
10	cctccgggtct cgacggaatc aactgctaag ttagggatca ctgggtgtag atctgatgcc	120
	aatcgagttt ttgccactgc tactgcgcgc gctacagcta cagctaccac cggtgagatt	180
	tcgtctagag ttgcggcttt ggctggatta gggcatcact acgctcggtg ttattgggag	240
	ctttctaaag ctaaacttag tatgcttggt gttgcaactt ctggaactgg gtatattctg	300
15	ggtacgggaa atgctgcaat tagcttcccg gggctttgtt acacatgtgc aggaaccatg	360
	atgattgctg catctgctaa ttccttgaat cagatttttg agataagcaa tgattctaag	420
	atgaaaagaa cgatgctaag gccattgcct tcaggacgta ttagtggtcc acacgctggt	480
	gcattgggcta ctattgctgg tgcttctgggt gcttgtttgt tggccagcaa gactaatatg	540
	ttggctgctg gaattgcac tgccaatcct gtactttatg cgtttgttta tactccgttg	600
20	aagcaacttc accctatcaa tacatggggt ggcgctgttg ttgggtgctat cccacccttg	660
	cttgggtggg cggcagcgtc tggtcagatt tcatacaatt cgatgattct tccagctgct	720
	ctttactttt ggcagatacc tcattttatg gcccttgac atctctgccg caatgattat	780
	gcagctggag gttacaagat gttgtcactc tttgatccgt caggaagag aatagcagca	840
	gtggctctaa ggaactgctt ttacatgac cctctcggtt tcatcgcta tgactggggg	900
25	ttaacctcaa gttggttttg cctcgaatca acactttctca cactagcaat cgctgcaaca	960
	gcattttcat tctaccgaga ccggaccatg cataaagcaa ggaaaatgtt ccatgccagt	1020
	cttctcttcc ttctgtttt catgtctggt cttctctac accgtgtctc taatgataat	1080
	cagcaacaac tcgtagaaga agccggatta acaaattctg tatctggtga agtcaaaact	1140
	cagaggcgaa agaaacgtgt ggctcaacct ccggtggctt atgcctctgc tgcaccgttt	1200
30	cctttcctcc cagctccttc cttctactct ccatga	1260
		1296

&lt;210&gt; 6

&lt;211&gt; 431

&lt;212&gt; PRT

35 &lt;213&gt; Arabidopsis sp

&lt;400&gt; 6

Met Trp Arg Arg Ser Val Val Tyr Arg Phe Ser Ser Arg Ile Ser Val

1

5

10

15

Ser Ser Ser Leu Pro Asn Pro Arg Leu Ile Pro Trp Ser Arg Glu Leu  
 20 25 30  
 Cys Ala Val Asn Ser Phe Ser Gln Pro Pro Val Ser Thr Glu Ser Thr  
 35 40 45  
 5 Ala Lys Leu Gly Ile Thr Gly Val Arg Ser Asp Ala Asn Arg Val Phe  
 50 55 60  
 Ala Thr Ala Thr Ala Ala Ala Thr Ala Thr Ala Thr Thr Gly Glu Ile  
 65 70 75 80  
 Ser Ser Arg Val Ala Ala Leu Ala Gly Leu Gly His His Tyr Ala Arg  
 10 85 90 95  
 Cys Tyr Trp Glu Leu Ser Lys Ala Lys Leu Ser Met Leu Val Val Ala  
 100 105 110  
 Thr Ser Gly Thr Gly Tyr Ile Leu Gly Thr Gly Asn Ala Ala Ile Ser  
 115 120 125  
 15 Phe Pro Gly Leu Cys Tyr Thr Cys Ala Gly Thr Met Met Ile Ala Ala  
 130 135 140  
 Ser Ala Asn Ser Leu Asn Gln Ile Phe Glu Ile Ser Asn Asp Ser Lys  
 145 150 155 160  
 Met Lys Arg Thr Met Leu Arg Pro Leu Pro Ser Gly Arg Ile Ser Val  
 20 165 170 175  
 Pro His Ala Val Ala Trp Ala Thr Ile Ala Gly Ala Ser Gly Ala Cys  
 180 185 190  
 Leu Leu Ala Ser Lys Thr Asn Met Leu Ala Ala Gly Leu Ala Ser Ala  
 195 200 205  
 25 Asn Leu Val Leu Tyr Ala Phe Val Tyr Thr Pro Leu Lys Gln Leu His  
 210 215 220  
 Pro Ile Asn Thr Trp Val Gly Ala Val Val Gly Ala Ile Pro Pro Leu  
 225 230 235 240  
 Leu Gly Trp Ala Ala Ala Ser Gly Gln Ile Ser Tyr Asn Ser Met Ile  
 30 245 250 255  
 Leu Pro Ala Ala Leu Tyr Phe Trp Gln Ile Pro His Phe Met Ala Leu  
 260 265 270  
 Ala His Leu Cys Arg Asn Asp Tyr Ala Ala Gly Gly Tyr Lys Met Leu  
 275 280 285  
 35 Ser Leu Phe Asp Pro Ser Gly Lys Arg Ile Ala Ala Val Ala Leu Arg  
 290 295 300  
 Asn Cys Phe Tyr Met Ile Pro Leu Gly Phe Ile Ala Tyr Asp Trp Gly  
 305 310 315 320  
 Leu Thr Ser Ser Trp Phe Cys Leu Glu Ser Thr Leu Leu Thr Leu Ala

325 330 335  
 Ile Ala Ala Thr Ala Phe Ser Phe Tyr Arg Asp Arg Thr Met His Lys  
 340 345 350  
 Ala Arg Lys Met Phe His Ala Ser Leu Leu Phe Leu Pro Val Phe Met  
 5 355 360 365  
 Ser Gly Leu Leu Leu His Arg Val Ser Asn Asp Asn Gln Gln Gln Leu  
 370 375 380  
 Val Glu Glu Ala Gly Leu Thr Asn Ser Val Ser Gly Glu Val Lys Thr  
 385 390 395 400  
 10 Gln Arg Arg Lys Lys Arg Val Ala Gln Pro Pro Val Ala Tyr Ala Ser  
 405 410 415  
 Ala Ala Pro Phe Pro Phe Leu Pro Ala Pro Ser Phe Tyr Ser Pro  
 420 425 430

15 <210> 7  
 <211> 479  
 <212> DNA  
 <213> Arabidopsis sp

20 <400> 7  
 ggaaactccc ggagcacctg ttgacaggta ccgctaacct taatcgataa tttattttctc 60  
 ttgtcaggaa ttatgtaagt ctggtggaag gctcgcatat catttttgca ttgcctttcg 120  
 ctatgatcgg gtttactttg ggtgtgatga gaccaggcgt ggcttttatgg tatggcgaaa 180  
 acccattttt atccaatgct gcattccctc ccgatgattc gttctttcat tcctatacag 240  
 25 gtatcatgct gataaaactg ttactggtac tggtttgtat ggtatcagca agaagcgagg 300  
 cgatggcggt taaccgggtat ctgcacaggc attttgacgc gaagaaccgc cgtactgcc 360  
 tccgtgaaat acctgcgggc gtcatatctg ccaacagtgc gctggtgttt acgataggct 420  
 gctgcgtggt attctgggtg gcctgttatt tcattaacac gatctgtttt tacctggcg 479

30 <210> 8  
 <211> 551  
 <212> DNA  
 <213> Arabidopsis sp

35 <220>  
 <221> misc\_feature  
 <222> (1) ... (551)  
 <223> n = A,T,C or G



<400> 8

	ttgtggctta caccttaatg agcatacgcc agnccattac ggctcgtaa tcggcgccat	60
	ngccgngct gntgcaccgg tagtgggcta ctgcgcctg accaatcagc ttgatctagc	120
	ggctcttatt ctgtttttaa ttttactgtt ctggcaaagc ccgcattttt acgcgatttc	180
5	cattttcagg ctaaaagact tttcagcggc ctgtattccg gtgctgcca tcattaaaga	240
	cctgcgctat accaaaatca gcatgctggt ttacgtgggc ttatttacac tggetgctat	300
	catgccggcc ctcttagggt atgccggtg gatttatggg atagcggcct taattttagg	360
	cttgtattgg ctttatattg ccatacaagg attcaagacc gccgatgatc aaaaatggtc	420
	tcgtaagatg tttggatctt cgattttaat cattaccctc ttgtcggtaa tgatgcttgt	480
10	ttaaacttac tgctcctga agtttatata tcgataattt cagcttaagg aggcttagtg	540
	gttaattcaa t	551

<210> 9

<211> 297

15 <212> PRT

<213> Arabidopsis sp

<400> 9

	Met Val Leu Ala Glu Val Pro Lys Leu Ala Ser Ala Ala Glu Tyr Phe	
20	1 5 10 15	
	Phe Lys Arg Gly Val Gln Gly Lys Gln Phe Arg Ser Thr Ile Leu Leu	
	20 25 30	
	Leu Met Ala Thr Ala Leu Asn Val Arg Val Pro Glu Ala Leu Ile Gly	
	35 40 45	
25	Glu Ser Thr Asp Ile Val Thr Ser Glu Leu Arg Val Arg Gln Arg Gly	
	50 55 60	
	Ile Ala Glu Ile Thr Glu Met Ile His Val Ala Ser Leu Leu His Asp	
	65 70 75 80	
	Asp Val Leu Asp Asp Ala Asp Thr Arg Arg Gly Val Gly Ser Leu Asn	
30	85 90 95	
	Val Val Met Gly Asn Lys Val Val Ala Leu Leu Ala Thr Ala Val Glu	
	100 105 110	
	His Leu Val Thr Gly Glu Thr Met Glu Ile Thr Ser Ser Thr Glu Gln	
	115 120 125	
35	Arg Tyr Ser Met Asp Tyr Tyr Met Gln Lys Thr Tyr Tyr Lys Thr Ala	
	130 135 140	
	Ser Leu Ile Ser Asn Ser Cys Lys Ala Val Ala Val Leu Thr Gly Gln	
	145 150 155 160	
	Thr Ala Glu Val Ala Val Leu Ala Phe Glu Tyr Gly Arg Asn Leu Gly	

	165	170	175
	Leu Ala Phe Gln Leu Ile Asp Asp Ile Leu Asp Phe Thr Gly Thr Ser		
	180	185	190
	Ala Ser Leu Gly Lys Gly Ser Leu Ser Asp Ile Arg His Gly Val Ile		
5	195	200	205
	Thr Ala Pro Ile Leu Phe Ala Met Glu Glu Phe Pro Gln Leu Arg Glu		
	210	215	220
	Val Val Asp Gln Val Glu Lys Asp Pro Arg Asn Val Asp Ile Ala Leu		
	225	230	235
10	Glu Tyr Leu Gly Lys Ser Lys Gly Ile Gln Arg Ala Arg Glu Leu Ala		240
	245	250	255
	Met Glu His Ala Asn Leu Ala Ala Ala Ala Ile Gly Ser Leu Pro Glu		
	260	265	270
	Thr Asp Asn Glu Asp Val Lys Arg Ser Arg Arg Ala Leu Ile Asp Leu		
15	275	280	285
	Thr His Arg Val Ile Thr Arg Asn Lys		
	290	295	
	<210> 10		
20	<211> 561		
	<212> DNA		
	<213> Arabidopsis sp		
	<400> 10		
25	aagcgcatcc gtcctcttct acgattgccg ccagccgcat gtatggctgc ataaccgacc		60
	gccccatatec gctcgcggcc gcggtcgaat tcattcacac cgcgacgctg ctgcatgacg		120
	acgtcgtcga tgaaagcgat ttgcgcgcgc gccgcgaaag cgcgcataag gttttcggca		180
	atcaggcgag cgtgctcgtc ggcgatttcc ttttctcccg cgccttcag ctgatggtgg		240
	aagacggctc gctcgacgcg ctgcgcattc tctcggatgc ctccgccgtg atcgcgcagg		300
30	gcgaagtgat gcagctcggc accgcgcgca atcttgaaac caatatgagc cagtatctcg		360
	atgtgatcag cgcaagacc gccgcgctct ttgcgcgcgc ctgcgaaatc ggcccgggtga		420
	tggcgaacgc gaaggcggaa gatgctgccg cgatgtgcga atacggcatg aatctcggta		480
	tgccttcca gatcatcgac gaccttctcg attacggcac cggcggccac gccgagcttg		540
	gcaagaacac gggcgacgat t		561
35	<210> 11		
	<211> 966		
	<212> DNA		
	<213> Arabidopsis sp		

<400> 11  
 atggtacttg ccgaggttcc aaagcttgcc tctgctgctg agtacttctt caaaaggggt 60  
 gtgcaaggaa aacagtttcg ttcaactatt ttgctgctga tggcgacagc tctgaatgta 120  
 5 cgcgttccag aagcattgat tggggaatca acagatatag tcacatcaga attacgcgta 180  
 aggcaacggg gtattgctga aatcactgaa atgatacacg tcgcaagtct actgcacgat 240  
 gatgtcttgg atgatgccga tacaaggcgt ggtgttggtt ccttaaagt tghtaatgggt 300  
 aacaagatgt cgggtattagc aggagacttc ttgctctccc gggcttgtgg ggctctcget 360  
 gcttttaaaga acacagaggt tgtagcatta cttgcaactg ctgtagaaca tcttgttacc 420  
 10 ggtgaaacca tggaaataac tagttcaacc gagcagcgtt atagtatgga ctactacatg 480  
 cagaagacat attataagac agcatcgcta atctctaaca gctgcaaagc tgttgccgtt 540  
 ctcactggac aaacagcaga agttgccgtg ttagcttttg agtatgggag gaatctgggt 600  
 ttagcattcc aattaataga cgacattctt gatttcacgg gcacatctgc ctctctcgga 660  
 aagggatcgt tgtcagatat tcgccatgga gtcataacag cccaatcct ctttgccatg 720  
 15 gaagagtttc ctcaactacg cgaagtgtt gatcaagttg aaaaagatcc taggaatgtt 780  
 gacattgctt tagagtatct tgggaagagc aagggaatac agagggaag agaattagcc 840  
 atggaacatg cgaatctagc agcagctgca atcggtctc tacctgaaac agacaatgaa 900  
 gatgtcaaaa gatcgaggcg ggcacttatt gacttgaccc atagagtcac caccagaac 960  
 aagtga 966

20

&lt;210&gt; 12

&lt;211&gt; 321

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis sp

25

&lt;400&gt; 12

Met Val Leu Ala Glu Val Pro Lys Leu Ala Ser Ala Ala Glu Tyr Phe

1 5 10 15

Phe Lys Arg Gly Val Gln Gly Lys Gln Phe Arg Ser Thr Ile Leu Leu

30 20 25 30

Leu Met Ala Thr Ala Leu Asn Val Arg Val Pro Glu Ala Leu Ile Gly

35 40 45

Glu Ser Thr Asp Ile Val Thr Ser Glu Leu Arg Val Arg Gln Arg Gly

50 55 60

35 Ile Ala Glu Ile Thr Glu Met Ile His Val Ala Ser Leu Leu His Asp

65 70 75 80

Asp Val Leu Asp Asp Ala Asp Thr Arg Arg Gly Val Gly Ser Leu Asn

85 90 95

Val Val Met Gly Asn Lys Met Ser Val Leu Ala Gly Asp Phe Leu Leu

100 105 110  
 Ser Arg Ala Cys Gly Ala Leu Ala Ala Leu Lys Asn Thr Glu Val Val  
 115 120 125  
 Ala Leu Leu Ala Thr Ala Val Glu His Leu Val Thr Gly Glu Thr Met  
 5 130 135 140  
 Glu Ile Thr Ser Ser Thr Glu Gln Arg Tyr Ser Met Asp Tyr Tyr Met  
 145 150 155 160  
 Gln Lys Thr Tyr Tyr Lys Thr Ala Ser Leu Ile Ser Asn Ser Cys Lys  
 165 170 175  
 10 Ala Val Ala Val Leu Thr Gly Gln Thr Ala Glu Val Ala Val Leu Ala  
 180 185 190  
 Phe Glu Tyr Gly Arg Asn Leu Gly Leu Ala Phe Gln Leu Ile Asp Asp  
 195 200 205  
 Ile Leu Asp Phe Thr Gly Thr Ser Ala Ser Leu Gly Lys Gly Ser Leu  
 15 210 215 220  
 Ser Asp Ile Arg His Gly Val Ile Thr Ala Pro Ile Leu Phe Ala Met  
 225 230 235 240  
 Glu Glu Phe Pro Gln Leu Arg Glu Val Val Asp Gln Val Glu Lys Asp  
 245 250 255  
 20 Pro Arg Asn Val Asp Ile Ala Leu Glu Tyr Leu Gly Lys Ser Lys Gly  
 260 265 270  
 Ile Gln Arg Ala Arg Glu Leu Ala Met Glu His Ala Asn Leu Ala Ala  
 275 280 285  
 Ala Ala Ile Gly Ser Leu Pro Glu Thr Asp Asn Glu Asp Val Lys Arg  
 25 290 295 300  
 Ser Arg Arg Ala Leu Ile Asp Leu Thr His Arg Val Ile Thr Arg Asn  
 305 310 315 320  
 Lys  
 30  
 <210> 13  
 <211> 621  
 <212> DNA  
 <213> Arabidopsis sp  
 35  
 <400> 13  
 gctttctcct ttgctaattc ttgagctttc ttgatccac cgcgatttct aactatttca 60  
 atcgtctctt caagcgatcc aggcacaaa aactcagact caatgatctc tcttagcctt 120  
 ggctcattct ctagcgcgaa gatcactggc gccgttatgt tacctttggc taagtcatta 180

	gctgcaggct tacctaactg ctctgtggac tgagtgaagt ccagaatgtc atcaactact	240
	tgaaaagata aaccgagatt cttcccgaac tgatacatTT gctctgcgac cttgctttcg	300
	actttactga aaattgctgc tcctttggtg cttgcagcta ctaatgaagc tgtctttag	360
	taactcttta gcatgtagtc atcaagcttg acatcacaat cgaataaact cgatgcttgc	420
5	tttatctcac cgcttgcaaa atctttgatc acctgcaaaa agataaatca agattcagac	480
	caaatgttct ttgtattgag tagcttcac taatctcaga aaggaaattt acctgactta	540
	tgagcttaat gacttcaagg ttttcgagat ttgtaagtac catgatgctt gagcaacatg	600
	aaatccccag ctaatacagc t	621
10	<210> 14	
	<211> 741	
	<212> DNA	
	<213> Arabidopsis sp	
15	<400> 14	
	ggtgagtttt gttaatagtt atgagattca tctatTTTTg tcataaaatt gtttggtttg	60
	gtttaaactc tgtgtataat tgcaggaaag gaaacagttc atgagctttt cggcacaaga	120
	gtagcgggtgc tagctggaga tttcatgttt gctcaagcgt catggtactt agcaaatctc	180
	gagaatcttg aagttattaa gctcatcagt cagggtactta gttactctta cattgttttt	240
20	ctatgaggtt gagctatgaa tctcatttcg ttgaataatg ctgtgcctca aacttttttt	300
	catgtttttca ggtgatcaaa gactttgcaa gcggagagat aaagcaggcg tccagcttat	360
	ttgactgcga caccaagctc gacgagtact tactcaaaag tttctacaag acagcctctt	420
	tagtgggtgc gagcaccaaa ggagctgcc a ttttcagcag agttgagcct gatgtgacag	480
	aacaaatgta cgagtttggg aagaatctcg gtctctcttt ccagatagtt gatgatattt	540
25	tggatttcac tcagtcgaca gacgagctcg ggaagccagc agggagtgat ttggctaaag	600
	gtaacttaac agcacctgtg attttcgctc tggagaggga gccaaaggcta agagagatca	660
	ttgagtcaaa gttctgtgag gcgggttctc ttggaagaagc gattgaagcg gtgacaaaag	720
	gtggggggat taagagagca c	741
30	<210> 15	
	<211> 1087	
	<212> DNA	
	<213> Arabidopsis sp	
35	<400> 15	
	cctcttcagc caatccagag gaagaagaga caacttttta tctttcgtca agagtctccg	60
	aaaacgcacg gttttatgct ctctcttctg ccctcacctc acaagacgca gggcacatga	120
	ttcaaccaga gggaaaaagc aacgataaca actctgcttt tgatttcaag ctgtatatga	180
	tccgcaaagc cgagtctgta aatgcggctc tcgacgtttc cgtaccgctt ctgaaacccc	240

	ttacgatcca agaagcgggc aggtactctt tgctagccgg cggaaaacgt gtgaggcctc	300
	tgctctgcat tgccgcttgt gagcttgtgg gggcgacga ggetactgcc atgtcagccg	360
	cttgccgggt cgagatgac cacacaagct ctctcattca tgacgatctt ccgtgcatgg	420
	acaatgccga cctccgtaga ggcaagccca ccaatcaca ggtatgttgt ttaattatat	480
5	gaaggctcag agataatgct gaactagtgt tgaaccaatt tttgctcaaa caaggtatat	540
	ggagaagaca tggcggtttt ggcaggtgat gcaactcctg cattggcgtt tgagcacatg	600
	acggttgtgt cgagtgggtt ggtcgctccc gagaagatga ttcgcgccgt ggttgagctg	660
	gccagggcca tagggactac agggctagtt gctggacaaa tgatagacct agccagcgaa	720
	agactgaatc cagacaaggt tggattggag catctagagt tcatccatct ccacaaaacg	780
10	gcggcattgt tggaggcagc ggcagtttta ggggttataa tgggaggtgg aacagaggaa	840
	gaaatcgaaa agcttagaaa gtatgctagg tgtattggac tactgtttca ggttgttgat	900
	gacattctcg acgtaacaaa atctactgag gaattgggta agacagccgg aaaagacgta	960
	atggccggaa agctgacgta tccaaggctg ataggtttgg agggatccag ggaagttgca	1020
	gagcacctga ggagagaagc agaggaaaag cttaaagggt ttgatccaag tcaggcggcg	1080
15	cctctgg	1087
	<210> 16	
	<211> 1164	
	<212> DNA	
20	<213> Arabidopsis sp	
	<400> 16	
	atgacttoga ttctcaacac tgtctccacc atccactctt ccagagttac ctccgtcgat	60
	cgagtcggag tctctctct tccgaattcg gattccgttg agttcactcg ccggcgcttct	120
25	ggtttctcga cgttgatcta cgaatcacc gggcgagat ttgttgtgcg tgcggcggag	180
	actgatactg ataaagttaa atctcagaca cctgacaagg caccagccgg tggttcaagc	240
	attaaccagc ttctcggtat caaaggagca tctcaagaaa ctaataaatg gaagattcgt	300
	cttcagctta caaaaccagt cacttggcct ccaactgggtt ggggagtcgt ctgtggtgct	360
	gctgcttcag ggaactttca ttggacccca gaggatgttg ctaagtcgat tctttgcatg	420
30	atgatgtctg gtcttgtct tactggctat acacagacaa tcaacgactg gtatgataga	480
	gatatcgacg caattaatga gccatatcgt ccaattccat ctggagcaat atcagagcca	540
	gaggttatta cacaagtctg ggtgctatta ttgggaggtc ttggtattgc tggaatatta	600
	gatgtgtggg cagggcatac cactcccact gtcttctatc ttgctttggg aggatcattg	660
	ctatcttata tatactctgc tccacctctt aagctaaaac aaaatggatg ggttgaaat	720
35	tttgcaattg gagcaagcta tattagtttg ccatggtggg ctggccaagc attgtttggc	780
	actcttacgc cagatgttgt tgttctaaca ctctgttaca gcatagctgg gttaggaata	840
	gccattgtta acgacttcaa aagtgttgaa ggagatagag cattaggact tcagtctctc	900
	ccagtagctt ttggcaccga aactgcaaaa tggatatgcy ttggtgctat agacattact	960
	cagctttctg ttgccggata tctattagca tctgggaaac cttattatgc gttggcgttg	1020

gttgetttga tcattcctca gattgtgttc cagtttaaat actttctcaa ggacctgtc 1080  
 aaatacgacg tcaagtacca ggcaagcgcg cagccattct tgggtgctcg aatatttgta 1140  
 acggcattag catcgcaaca ctga 1164

5 <210> 17  
 <211> 387  
 <212> PRT  
 <213> Arabidopsis sp

10 <400> 17  
 Met Thr Ser Ile Leu Asn Thr Val Ser Thr Ile His Ser Ser Arg Val  
 1 5 10 15  
 Thr Ser Val Asp Arg Val Gly Val Leu Ser Leu Arg Asn Ser Asp Ser  
 20 25 30

15 Val Glu Phe Thr Arg Arg Arg Ser Gly Phe Ser Thr Leu Ile Tyr Glu  
 35 40 45  
 Ser Pro Gly Arg Arg Phe Val Val Arg Ala Ala Glu Thr Asp Thr Asp  
 50 55 60  
 Lys Val Lys Ser Gln Thr Pro Asp Lys Ala Pro Ala Gly Gly Ser Ser  
 65 70 75 80  
 Ile Asn Gln Leu Leu Gly Ile Lys Gly Ala Ser Gln Glu Thr Asn Lys  
 85 90 95  
 Trp Lys Ile Arg Leu Gln Leu Thr Lys Pro Val Thr Trp Pro Pro Leu  
 100 105 110

25 Val Trp Gly Val Val Cys Gly Ala Ala Ala Ser Gly Asn Phe His Trp  
 115 120 125  
 Thr Pro Glu Asp Val Ala Lys Ser Ile Leu Cys Met Met Met Ser Gly  
 130 135 140  
 Pro Cys Leu Thr Gly Tyr Thr Gln Thr Ile Asn Asp Trp Tyr Asp Arg  
 145 150 155 160  
 Asp Ile Asp Ala Ile Asn Glu Pro Tyr Arg Pro Ile Pro Ser Gly Ala  
 165 170 175  
 Ile Ser Glu Pro Glu Val Ile Thr Gln Val Trp Val Leu Leu Leu Gly  
 180 185 190

35 Gly Leu Gly Ile Ala Gly Ile Leu Asp Val Trp Ala Gly His Thr Thr  
 195 200 205  
 Pro Thr Val Phe Tyr Leu Ala Leu Gly Gly Ser Leu Leu Ser Tyr Ile  
 210 215 220  
 Tyr Ser Ala Pro Pro Leu Lys Leu Lys Gln Asn Gly Trp Val Gly Asn

	<210>	18	
	<211>	981	
25	<212>	DNA	
	<213>	Arabidopsis sp	
	<400>	18	
30	atgttgttta	gtggttcagc	gatcccatta agcagcttct gctctcttcc ggagaaaccc 60
	cacactcttc	ctatgaaact	ctctcccgcct gcaatccgat cttcatcctc atctgccccg 120
	gggtcgttga	acttcgatct	gaggacgtat tggacgactc tgatcaccga gatcaaccag 180
	aagctggatg	aggccatacc	ggtcaagcac cctgcgggga tctacgaggc tatgagatac 240
	tctgtactcg	cacaaggcgc	caagcgtgcc cctcctgtga tgtgtgtggc ggcttgcgag 300
	ctcttcggtg	gcgatcgctt	cgccgctttc cccaccgcct gtgccctaga aatggtgcac 360
35	gcggtcttct	tgatacacga	cgacctcccc tgtatggacg acgatcctgt gcgcagagga 420
	aagccatcta	accacactgt	ctacggctct ggcatggcca ttctcgccgg tgacgcctc 480
	ttcccactcg	ccttcagca	cattgtctcc cacacgcctc ctgaccttgt tccccgagcc 540
	accatctcta	gactcatcac	tgagattgcc cgactgtcg gctccactgg tatggctgca 600
	ggccagtagc	tcgaccttga	aggaggcccc ttctctcttt cctttgttca ggagaagaaa 660



	ttcggagcca tgggtgaatg ctctgccgtg tgcggtggcc tattgggcgg tgccactgag	720
	gatgagctcc agagtctccg aaggtaacggg agagccgtcg ggatgctgta tcaggtggtc	780
	gatgacatca ccgaggacaa gaagaagagc tatgatgggtg gagcagagaa gggaatgatg	840
	gaaatggcgg aagagctcaa ggagaaggcg aagaaggagc ttcaagtgtt tgacaacaag	900
5	tatggaggag gagacacact tgttcctctc tacaccttcg ttgactacgc tgctcatcga	960
	cattttcttc ttcccctctg a	981
	<210> 19	
	<211> 245	
10	<212> DNA	
	<213> GLycine sp	
	<400> 19	
	gcaacatctg ggactgggtt tgtcttgggg agtggtagtg ctgttgatct ttcggcactt	60
15	tcttgcaact gcttgggtac catgatgggt gctgcactcg ctaactcttt gaatcagggt	120
	tttgagatca ataatgatgc taaaatgaag agaacaagtc gcaggecact accctcagga	180
	cgcatacaaa tacctcatgc agttggctgg gcatcctctg ttggattagc tggtaacggct	240
	ctact	245
20	<210> 20	
	<211> 253	
	<212> DNA	
	<213> Glycine sp	
25	<400> 20	
	attggctttc caagatcatt gggttttctt gttgcattca tgaccttcta ctcttgggt	60
	ttggcattgt ccaaggatat acctgacgtt gaaggagata aagagcacgg cattgattct	120
	tttgagtagc gtctaggtca gaaacgggca ttttggattt gcgtttcctt ttttgaaatg	180
	gctttcggag ttggtatcct ggccggagca tcatgctcac acttttggac taaaattttc	240
30	acgggtatgg gaa	253
	<210> 21	
	<211> 275	
	<212> DNA	
35	<213> Glycine sp	
	<400> 21	
	tgatcttcta ctctctgggt atggcattgt ccaaggatat atctgacgtt aaaggagata	60
	aagcatacgg catcgatact ttagcgatac gtttgggtca aaaatgggta ttttggattt	120

	gcattatcct ttttgaaatg gcttttggag ttgccctctt ggcaggagca acatcttctt	180
	acctttggat taaaattgtc acgggtctgg gacatgctat tcttgettca attctcttgt	240
	accaagccaa atctatatac ttgagcaaca aagtt	275
5	<210> 22	
	<211> 299	
	<212> DNA	
	<213> Glycine sp	
10	<220>	
	<221> misc_feature	
	<222> (1) ... (299)	
	<223> n = A,T,C or G	
15	<400> 22	
	ccanaatang tncatcttng aaagacaatt ggcctcttca acacacaagt ctgcatgtga	60
	agaagaggcc aattgtcttt ccaagatcac ttatngtggc tattgtaatc atgaacttct	120
	tctttgtggg tatggcattg gcaaaggata tacctantcg ttgaaggaga taaaatatat	180
	ggcattgata cttttgcaat acgtataggt caaaaacaag tattttggat ttgtattttc	240
20	ctttttgaaa ggctttcgga gtttccttag tggcaggagc aacatcttct agccttggg	299
	<210> 23	
	<211> 767	
	<212> DNA	
25	<213> Glycine sp	
	<400> 23	
	gtggaggctg tggttgctgc cctgtttatg aatatttata ttgttggttt gaatcaattg	60
	tctgatgttg aaatagacaa gataaacaag ccgtatcttc cattagcacc tggggaatat	120
30	tcctttgaaa ctggtgtcac tattgttgca tctttttcaa ttctgagttt ttggcttggc	180
	tgggtttag gttcatggcc attattttgg gccctttttg taagctttgt gctaggaact	240
	gcttattcaa tcaatgtgcc tctgttgaga tgggaagagg ttgcagtgc tgcagcgatg	300
	tgcattctag ctgttcgggc agtaatagtt caacttgcac ttttccttca catgcagact	360
	catgtgtaca agaggccacc tgtcttttca agaccattga tttttgctac tgcattcatg	420
35	agcttcttct ctgtagttat agcactgttt aaggatatac ctgacattga aggagataaa	480
	gtatttggca tccaatcttt ttcagtgtgt ttaggtcaga agccggtgtt ctggacttgt	540
	gttacccttc ttgaaatagc ttatggagtc gccctcctgg tgggagctgc atctccttgt	600
	ctttggagca aaattttcac gggctctggga cacgctgtgc tggcttcaat tctctggttt	660
	catgccaaat ctgtagattt gaaaagcaaa gcttcgataa catccttcta tatgtttatt	720

tggaagctat tttatgcaga atacttactc attccttttg ttagatg

767

<210> 24

<211> 255

5 <212> PRT

<213> Glycine sp

<400> 24

Val Glu Ala Val Val Ala Ala Leu Phe Met Asn Ile Tyr Ile Val Gly  
 10 1 5 10 15  
 Leu Asn Gln Leu Ser Asp Val Glu Ile Asp Lys Ile Asn Lys Pro Tyr  
 20 25 30  
 Leu Pro Leu Ala Ser Gly Glu Tyr Ser Phe Glu Thr Gly Val Thr Ile  
 35 40 45  
 15 Val Ala Ser Phe Ser Ile Leu Ser Phe Trp Leu Gly Trp Val Val Gly  
 50 55 60  
 Ser Trp Pro Leu Phe Trp Ala Leu Phe Val Ser Phe Val Leu Gly Thr  
 65 70 75 80  
 Ala Tyr Ser Ile Asn Val Pro Leu Leu Arg Trp Lys Arg Phe Ala Val  
 20 85 90 95  
 Leu Ala Ala Met Cys Ile Leu Ala Val Arg Ala Val Ile Val Gln Leu  
 100 105 110  
 Ala Phe Phe Leu His Met Gln Thr His Val Tyr Lys Arg Pro Pro Val  
 115 120 125  
 25 Phe Ser Arg Pro Leu Ile Phe Ala Thr Ala Phe Met Ser Phe Phe Ser  
 130 135 140  
 Val Val Ile Ala Leu Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Lys  
 145 150 155 160  
 Val Phe Gly Ile Gln Ser Phe Ser Val Cys Leu Gly Gln Lys Pro Val  
 30 165 170 175  
 Phe Trp Thr Cys Val Thr Leu Leu Glu Ile Ala Tyr Gly Val Ala Leu  
 180 185 190  
 Leu Val Gly Ala Ala Ser Pro Cys Leu Trp Ser Lys Ile Phe Thr Gly  
 195 200 205  
 35 Leu Gly His Ala Val Leu Ala Ser Ile Leu Trp Phe His Ala Lys Ser  
 210 215 220  
 Val Asp Leu Lys Ser Lys Ala Ser Ile Thr Ser Phe Tyr Met Phe Ile  
 225 230 235 240  
 Trp Lys Leu Phe Tyr Ala Glu Tyr Leu Leu Ile Pro Phe Val Arg

245

250

255

<210> 25  
 <211> 360  
 5 <212> DNA  
 <213> Zea sp

<220>  
 <221> misc\_feature  
 10 <222> (1) ... (360)  
 <223> n = A,T,C or G

<400> 25  
 15 ggcgtcttca cttgttctgg tcttctogta tccctgatg aagaggttca cattttggcc 60  
 tcaggcttat cttggcctga cattcaactg gggagcttta ctagggtggg ctgctattaa 120  
 ggaaagcata gaccctgcaa atcatccttc cattgtatac agctgggtatt tgttggacgc 180  
 tgggtgatga tactatataat ggcacacagg tgtttcgcta tccctacttt catattaatc 240  
 cttgatgaag tggccatttc atgttgctgc ggtggtctta tacttgcata tctccatgca 300  
 tctcaggaca aagangatga cctgaaagta ggagtccaag tccacagctt aagatttggg 360

20  
 <210> 26  
 <211> 299  
 <212> DNA  
 <213> Zea sp

25  
 <220>  
 <221> misc\_feature  
 <222> (1) ... (299)  
 <223> n = A,T,C or G

30  
 <400> 26  
 gatggttgca gcactgcaa ataccctcaa ccagggtgtt gngataaaaa atgatgctaa 60  
 aatgaaaagg acaatgcgtg cccctgcca tctggtcgca ttagtctgc acatgetgcg 120  
 atgtgggcta caagtgttg agttgcagga acagctttgt tggcctggaa ggctaattggc 180  
 35 ttggcagctg ggcttgacgc ttctaatttt gttctgtatg catttgtgta tacgcggtg 240  
 aagcaaatac accctgttaa tacatgggtt ggggcagtcg ttggtgccat cccaccact 299

<210> 27  
 <211> 255

<212> DNA  
 <213> Zea sp

<220>

5 <221> misc\_feature  
 <222> (1) ... (255)  
 <223> n = A,T,C or G

<400> 27

10	anacttgc	atctccatgc	ntctcaggac	aaagangatg	acctgaaagt	aggtgtcaag	60
	tccacagcat	taagatttgg	agatttgacc	nnatactgna	tcagtggctt	tggcgcgga	120
	tgttcggca	gcttagcact	cagtggttac	aatgctgacc	ttggttggtg	tttagtgtga	180
	tgcttgagcg	aagaatggta	tngtttttac	ttgatattga	ctccagacct	gaaatcatgt	240
	tggaacaggt	ggccc					255

15

<210> 28  
 <211> 257  
 <212> DNA  
 <213> Zea sp

20

<400> 28

	attgaagggg	ataggactct	ggggcttcag	tcacttcctg	ttgcttttgg	gatggaaact	60
	gcaaaatgga	tttgtgttgg	agcaattgat	atcactcaat	tatctgttgc	aggttaccta	120
	ttgagcaccg	gtaagctgta	ttatgccctg	gtgttgcttg	ggctaacaat	tcctcaggtg	180
25	ttctttcagt	tccagtactt	cctgaaggac	cctgtgaagt	atgatgtcaa	atatcaggca	240
	agcgcacaa	cattctt					257

<210> 29  
 <211> 368

30 <212> DNA  
 <213> Zea sp

<400> 29

	atccagttgc	aaataataat	ggcgttcttc	tctgttgtaa	tagcactatt	caaggatata	60
35	cctgacatcg	aaggggaccg	catattcggg	atccgatcct	tcagcgccg	gttagggcaa	120
	aagaaggtct	tttgatctg	cggtggcttg	cttgagatgg	cctacagcgt	tgcgatactg	180
	atgggagcta	cctcttctg	tttgtggagc	aaaacagcaa	ccatcgctgg	ccattccata	240
	cttgccgcga	tcctatggag	ctgcgcgcga	tcggtggact	tgacgagcaa	agccgcaata	300
	acgtccttct	acatgttcat	ctggaagctg	ttctacgcgg	agtacctgct	catccctctg	360

gtgcggtg

368

&lt;210&gt; 30

&lt;211&gt; 122

5 &lt;212&gt; PRT

&lt;213&gt; Zea sp

&lt;400&gt; 30

Ile Gln Leu Gln Ile Ile Met Ala Phe Phe Ser Val Val Ile Ala Leu  
 10 1 5 10 15  
 Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Arg Ile Phe Gly Ile Arg  
 20 25 30  
 Ser Phe Ser Val Arg Leu Gly Gln Lys Lys Val Phe Trp Ile Cys Val  
 35 40 45  
 15 Gly Leu Leu Glu Met Ala Tyr Ser Val Ala Ile Leu Met Gly Ala Thr  
 50 55 60  
 Ser Ser Cys Leu Trp Ser Lys Thr Ala Thr Ile Ala Gly His Ser Ile  
 65 70 75 80  
 Leu Ala Ala Ile Leu Trp Ser Cys Ala Arg Ser Val Asp Leu Thr Ser  
 20 85 90 95  
 Lys Ala Ala Ile Thr Ser Phe Tyr Met Phe Ile Trp Lys Leu Phe Tyr  
 100 105 110  
 Ala Glu Tyr Leu Leu Ile Pro Leu Val Arg  
 115 120

25

&lt;210&gt; 31

&lt;211&gt; 278

&lt;212&gt; DNA

&lt;213&gt; Zea sp

30

&lt;400&gt; 31

tattcagcac cacctctcaa gctcaagcag aatggatgga ttgggaactt cgctctgggt 60  
 gcgagttaca tcagcttgcc ctggtgggct ggccaggcgt tatttggaaac tcttacacca 120  
 gatattcattg tcttgactac tttgtacagc atagctgggc tagggattgc tattgtaaat 180  
 35 gatttcaaga gtattgaagg ggataggact ctggggcttc agtcacttcc tgttgctttt 240  
 gggatggaaa ctgcaaaatg gatttgtgtt ggagcaat 278

&lt;210&gt; 32

&lt;211&gt; 292

&lt;212&gt; PRT

&lt;213&gt; Synechocystis sp

&lt;400&gt; 32

5 Met Val Ala Gln Thr Pro Ser Ser Pro Pro Leu Trp Leu Thr Ile Ile  
     1                    5                    10                    15  
 Tyr Leu Leu Arg Trp His Lys Pro Ala Gly Arg Leu Ile Leu Met Ile  
                     20                    25                    30  
 Pro Ala Leu Trp Ala Val Cys Leu Ala Ala Gln Gly Leu Pro Pro Leu  
 10                    35                    40                    45  
 Pro Leu Leu Gly Thr Ile Ala Leu Gly Thr Leu Ala Thr Ser Gly Leu  
                     50                    55                    60  
 Gly Cys Val Val Asn Asp Leu Trp Asp Arg Asp Ile Asp Pro Gln Val  
                     65                    70                    75                    80  
 15 Glu Arg Thr Lys Gln Arg Pro Leu Ala Ala Arg Ala Leu Ser Val Gln  
                     85                    90                    95  
 Val Gly Ile Gly Val Ala Leu Val Ala Leu Leu Cys Ala Ala Gly Leu  
                     100                    105                    110  
 Ala Phe Tyr Leu Thr Pro Leu Ser Phe Trp Leu Cys Val Ala Ala Val  
 20                    115                    120                    125  
 Pro Val Ile Val Ala Tyr Pro Gly Ala Lys Arg Val Phe Pro Val Pro  
                     130                    135                    140  
 Gln Leu Val Leu Ser Ile Ala Trp Gly Phe Ala Val Leu Ile Ser Trp  
                     145                    150                    155                    160  
 25 Ser Ala Val Thr Gly Asp Leu Thr Asp Ala Thr Trp Val Leu Trp Gly  
                     165                    170                    175  
 Ala Thr Val Phe Trp Thr Leu Gly Phe Asp Thr Val Tyr Ala Met Ala  
                     180                    185                    190  
 Asp Arg Glu Asp Asp Arg Arg Ile Gly Val Asn Ser Ser Ala Leu Phe  
 30                    195                    200                    205  
 Phe Gly Gln Tyr Val Gly Glu Ala Val Gly Ile Phe Phe Ala Leu Thr  
                     210                    215                    220  
 Ile Gly Cys Leu Phe Tyr Leu Gly Met Ile Leu Met Leu Asn Pro Leu  
                     225                    230                    235                    240  
 35 Tyr Trp Leu Ser Leu Ala Ile Ala Ile Val Gly Trp Val Ile Gln Tyr  
                     245                    250                    255  
 Ile Gln Leu Ser Ala Pro Thr Pro Glu Pro Lys Leu Tyr Gly Gln Ile  
                     260                    265                    270  
 Phe Gly Gln Asn Val Ile Ile Gly Phe Val Leu Leu Ala Gly Met Leu

24



225                      230                      235                      240  
 Pro Leu His Gln Leu Gly Ile Leu Tyr Leu Ala Ile Ala Ile Ile Leu  
                                  245                      250                      255  
 Gly Gly Gln Phe Leu Val Lys Ala Trp Gln Leu Lys Gln Ala Pro Gly  
 5                      260                      265                      270  
 Asp Arg Asp Leu Ala Arg Gly Leu Phe Lys Phe Ser Ile Phe Tyr Leu  
                                  275                      280                      285  
 Met Leu Leu Cys Leu Ala Met Val Ile Asp Ser Leu Pro Val Thr His  
                                  290                      295                      300  
 10 Gln Leu Val Ala Gln Met Gly Thr Leu Leu Leu Gly  
                                  305                      310                      315  
  
 <210> 34  
 <211> 324  
 15 <212> PRT  
      <213> Synechocystis sp  
  
 <400> 34  
 Met Ser Asp Thr Gln Asn Thr Gly Gln Asn Gln Ala Lys Ala Arg Gln  
 20    1                      5                      10                      15  
 Leu Leu Gly Met Lys Gly Ala Ala Pro Gly Glu Ser Ser Ile Trp Lys  
                                  20                      25                      30  
 Ile Arg Leu Gln Leu Met Lys Pro Ile Thr Trp Ile Pro Leu Ile Trp  
                                  35                      40                      45  
 25 Gly Val Val Cys Gly Ala Ala Ser Ser Gly Gly Tyr Ile Trp Ser Val  
                                  50                      55                      60  
 Glu Asp Phe Leu Lys Ala Leu Thr Cys Met Leu Leu Ser Gly Pro Leu  
 65                      70                      75                      80  
 Met Thr Gly Tyr Thr Gln Thr Leu Asn Asp Phe Tyr Asp Arg Asp Ile  
 30                      85                      90                      95  
 Asp Ala Ile Asn Glu Pro Tyr Arg Pro Ile Pro Ser Gly Ala Ile Ser  
                                  100                      105                      110  
 Val Pro Gln Val Val Thr Gln Ile Leu Ile Leu Leu Val Ala Gly Ile  
                                  115                      120                      125  
 35 Gly Val Ala Tyr Gly Leu Asp Val Trp Ala Gln His Asp Phe Pro Ile  
                                  130                      135                      140  
 Met Met Val Leu Thr Leu Gly Gly Ala Phe Val Ala Tyr Ile Tyr Ser  
 145                      150                      155                      160  
 Ala Pro Pro Leu Lys Leu Lys Gln Asn Gly Trp Leu Gly Asn Tyr Ala

165 170 175  
 Leu Gly Ala Ser Tyr Ile Ala Leu Pro Trp Trp Ala Gly His Ala Leu  
 180 185 190  
 Phe Gly Thr Leu Asn Pro Thr Ile Met Val Leu Thr Leu Ile Tyr Ser  
 5 195 200 205  
 Leu Ala Gly Leu Gly Ile Ala Val Val Asn Asp Phe Lys Ser Val Glu  
 210 215 220  
 Gly Asp Arg Gln Leu Gly Leu Lys Ser Leu Pro Val Met Phe Gly Ile  
 225 230 235 240  
 10 Gly Thr Ala Ala Trp Ile Cys Val Ile Met Ile Asp Val Phe Gln Ala  
 245 250 255  
 Gly Ile Ala Gly Tyr Leu Ile Tyr Val His Gln Gln Leu Tyr Ala Thr  
 260 265 270  
 Ile Val Leu Leu Leu Leu Ile Pro Gln Ile Thr Phe Gln Asp Met Tyr  
 15 275 280 285  
 Phe Leu Arg Asn Pro Leu Glu Asn Asp Val Lys Tyr Gln Ala Ser Ala  
 290 295 300  
 Gln Pro Phe Leu Val Phe Gly Met Leu Ala Thr Gly Leu Ala Leu Gly  
 305 310 315 320  
 20 His Ala Gly Ile

&lt;210&gt; 35

&lt;211&gt; 307

25 &lt;212&gt; PRT

&lt;213&gt; Synechocystis sp

&lt;400&gt; 35

Met Thr Glu Ser Ser Pro Leu Ala Pro Ser Thr Ala Pro Ala Thr Arg  
 30 1 5 10 15  
 Lys Leu Trp Leu Ala Ala Ile Lys Pro Pro Met Tyr Thr Val Ala Val  
 20 25 30  
 Val Pro Ile Thr Val Gly Ser Ala Val Ala Tyr Gly Leu Thr Gly Gln  
 35 40 45  
 35 Trp His Gly Asp Val Phe Thr Ile Phe Leu Leu Ser Ala Ile Ala Ile  
 50 55 60  
 Ile Ala Trp Ile Asn Leu Ser Asn Asp Val Phe Asp Ser Asp Thr Gly  
 65 70 75 80  
 Ile Asp Val Arg Lys Ala His Ser Val Val Asn Leu Thr Gly Asn Arg

	85	90	95
	Asn Leu Val Phe Leu Ile Ser Asn Phe Phe Leu Leu Ala Gly Val Leu		
	100	105	110
	Gly Leu Met Ser Met Ser Trp Arg Ala Gln Asp Trp Thr Val Leu Glu		
5	115	120	125
	Leu Ile Gly Val Ala Ile Phe Leu Gly Tyr Thr Tyr Gln Gly Pro Pro		
	130	135	140
	Phe Arg Leu Gly Tyr Leu Gly Leu Gly Glu Leu Ile Cys Leu Ile Thr		
	145	150	155
10	Phe Gly Pro Leu Ala Ile Ala Ala Ala Tyr Tyr Ser Gln Ser Gln Ser		
	165	170	175
	Phe Ser Trp Asn Leu Leu Thr Pro Ser Val Phe Val Gly Ile Ser Thr		
	180	185	190
	Ala Ile Ile Leu Phe Cys Ser His Phe His Gln Val Glu Asp Asp Leu		
15	195	200	205
	Ala Ala Gly Lys Lys Ser Pro Ile Val Arg Leu Gly Thr Lys Leu Gly		
	210	215	220
	Ser Gln Val Leu Thr Leu Ser Val Val Ser Leu Tyr Leu Ile Thr Ala		
	225	230	235
20	Ile Gly Val Leu Cys His Gln Ala Pro Trp Gln Thr Leu Leu Ile Ile		
	245	250	255
	Ala Ser Leu Pro Trp Ala Val Gln Leu Ile Arg His Val Gly Gln Tyr		
	260	265	270
	His Asp Gln Pro Glu Gln Val Ser Asn Cys Lys Phe Ile Ala Val Asn		
25	275	280	285
	Leu His Phe Phe Ser Gly Met Leu Met Ala Ala Gly Tyr Gly Trp Ala		
	290	295	300
	Gly Leu Gly		
	305		
30			
	<210> 36		
	<211> 927		
	<212> DNA		
	<213> Synechocystis sp		
35			
	<400> 36		
	atggcaacta tccaagcttt ttggcgcttc tcccgcccc ataccatcat tggtaact	60	
	ctgagcgtct gggctgtgta tctgttaact attctcgggg atggaaactc agttaactcc	120	
	cctgcttccc tggatttagt gttcggcgct tggctggcct gcctgttggg taatgtgtac	180	

attgtcggcc tcaaccaatt gtgggatgtg gacattgacc gcatcaataa gccgaatttg 240  
 cccctagcta acggagattt ttctatcgcc cagggccggtt ggattgtggg actttgtgga 300  
 gttgcttcct tggcgatcgc ctggggatta gggctatggc tggggctaac ggtgggcatt 360  
 agtttgatta ttggcacggc ctattcgggtg ccgccagtga ggtaaagcg cttttccctg 420  
 5 ctggcggccc tgtgtattct gacggtgcgg ggaattgtgg ttaacttggg cttattttta 480  
 ttttttagaa ttggtttagg ttatcccccc actttaataa ccccatctg ggttttgact 540  
 ttatttatct tagttttcac cgtggcgatc gccattttta aagatgtgcc agatatggaa 600  
 ggcgatcggc aatttaagat tcaaacttta actttgcaa tggcaaaca aaacgttttt 660  
 cggggaacct taattttact cactggttgt tatntagcca tggcaatctg gggttatgg 720  
 10 gcggctatgc ctttaaatac tgctttcttg attgtttccc atttgtgctt attagcctta 780  
 ctctggtggc ggagtcgaga tgtacactta gaaagcaaaa ccgaaattgc tagtttttat 840  
 cagtttattt ggaagctatt tttcttagag tacttgctgt atcccttggc tctgtggtta 900  
 cctaattttt ctaatactat ttttttag 927

15 <210> 37  
 <211> 308  
 <212> PRT  
 <213> *Synechocystis* sp

20 <400> 37  
 Met Ala Thr Ile Gln Ala Phe Trp Arg Phe Ser Arg Pro His Thr Ile  
 1 5 10 15  
 Ile Gly Thr Thr Leu Ser Val Trp Ala Val Tyr Leu Leu Thr Ile Leu  
 20 25 30  
 25 Gly Asp Gly Asn Ser Val Asn Ser Pro Ala Ser Leu Asp Leu Val Phe  
 35 40 45  
 Gly Ala Trp Leu Ala Cys Leu Leu Gly Asn Val Tyr Ile Val Gly Leu  
 50 55 60  
 Asn Gln Leu Trp Asp Val Asp Ile Asp Arg Ile Asn Lys Pro Asn Leu  
 30 65 70 75 80  
 Pro Leu Ala Asn Gly Asp Phe Ser Ile Ala Gln Gly Arg Trp Ile Val  
 85 90 95  
 Gly Leu Cys Gly Val Ala Ser Leu Ala Ile Ala Trp Gly Leu Gly Leu  
 100 105 110  
 35 Trp Leu Gly Leu Thr Val Gly Ile Ser Leu Ile Ile Gly Thr Ala Tyr  
 115 120 125  
 Ser Val Pro Pro Val Arg Leu Lys Arg Phe Ser Leu Leu Ala Ala Leu  
 130 135 140  
 Cys Ile Leu Thr Val Arg Gly Ile Val Val Asn Leu Gly Leu Phe Leu

	145	150	155	160
	Phe Phe Arg Ile Gly Leu Gly Tyr Pro Pro Thr Leu Ile Thr Pro Ile			
		165	170	175
	Trp Val Leu Thr Leu Phe Ile Leu Val Phe Thr Val Ala Ile Ala Ile			
5		180	185	190
	Phe Lys Asp Val Pro Asp Met Glu Gly Asp Arg Gln Phe Lys Ile Gln			
		195	200	205
	Thr Leu Thr Leu Gln Ile Gly Lys Gln Asn Val Phe Arg Gly Thr Leu			
		210	215	220
10	Ile Leu Leu Thr Gly Cys Tyr Leu Ala Met Ala Ile Trp Gly Leu Trp			
		225	230	235
	Ala Ala Met Pro Leu Asn Thr Ala Phe Leu Ile Val Ser His Leu Cys			240
		245	250	255
	Leu Leu Ala Leu Leu Trp Trp Arg Ser Arg Asp Val His Leu Glu Ser			
15		260	265	270
	Lys Thr Glu Ile Ala Ser Phe Tyr Gln Phe Ile Trp Lys Leu Phe Phe			
		275	280	285
	Leu Glu Tyr Leu Leu Tyr Pro Leu Ala Leu Trp Leu Pro Asn Phe Ser			
		290	295	300
20	Asn Thr Ile Phe			
	305			
	<210> 38			
	<211> 1092			
25	<212> DNA			
	<213> Synechocystis sp			
	<400> 38			
	atgaaatttc cgccccacag tggttacat tggcaaggtc aatcaccttt ctttgaaggt			60
30	tggtagctgc gctgtctttt gccccaatcc ggggaaagtt ttgcttttat gtactccatc			120
	gaaaatcctg ctagcgatca tcattacggc ggcggtgctg tgcaaatttt agggccggct			180
	acgaaaaaac aagaaaatca ggaagaccaa cttgtttggc ggacatttcc ctcggtaaaa			240
	aaattttggg ccagtcctcg ccagtttgcc ctagggcatt ggggaaaatg tagggataac			300
	aggcaggcga aacccctact ctccgaagaa ttttttgcca cggcaagga aggttatcaa			360
35	atccatcaaa atcagcacca aggacaaatc attcatggcg atcgccattg tcgttggcag			420
	ttcacgtag aaccggaagt aacttggggg agtcctaacc gatttcctcg ggctacagcg			480
	ggttggcttt cctttttacc cttgtttgat cccggttggc aaattctttt agcccaaggt			540
	agagcgcacg gctggctgaa atggcagagg gaacagtatg aatttgacca cgccctagtt			600
	tatgccgaaa aaaattgggg tcactccttt ccctcccgct ggttttggct ccaagcaaat			660

tatttttctg accatccagg actgagcgtc actgccgctg gcggggaacg gattgttctt 720  
 ggtcgccccg aagaggtagc tttaattggc ttacatcacc aaggaattt ttacgaattt 780  
 ggcccgggcc atggcacagt cacttggcaa gtagctccct ggggccgttg gcaattaaaa 840  
 gccagcaatg ataggtattg ggtcaagttg tccggaaaaa cagataaaaa aggcagttta 900  
 5 gtccacactc ccaccgcca gggcttacia ctcaactgcc gagataccac taggggetat 960  
 ttgtatttgc aattgggacg tgtgggtcac ggcctgatag tgcaagggga aacggacacc 1020  
 gcggggctag aagttggagg tgattggggt ttaacagagg aaaatttgag caaaaaaaca 1080  
 gtgccattct ga 1092

10 <210> 39  
 <211> 363  
 <212> PRT  
 <213> *Synechocystis* sp

15 <400> 39  
 Met Lys Phe Pro Pro His Ser Gly Tyr His Trp Gln Gly Gln Ser Pro  
 1 5 10 15  
 Phe Phe Glu Gly Trp Tyr Val Arg Leu Leu Leu Pro Gln Ser Gly Glu  
 20 25 30  
 20 Ser Phe Ala Phe Met Tyr Ser Ile Glu Asn Pro Ala Ser Asp His His  
 35 40 45  
 Tyr Gly Gly Gly Ala Val Gln Ile Leu Gly Pro Ala Thr Lys Lys Gln  
 50 55 60  
 Glu Asn Gln Glu Asp Gln Leu Val Trp Arg Thr Phe Pro Ser Val Lys  
 25 65 70 75 80  
 Lys Phe Trp Ala Ser Pro Arg Gln Phe Ala Leu Gly His Trp Gly Lys  
 85 90 95  
 Cys Arg Asp Asn Arg Gln Ala Lys Pro Leu Leu Ser Glu Glu Phe Phe  
 100 105 110  
 30 Ala Thr Val Lys Glu Gly Tyr Gln Ile His Gln Asn Gln His Gln Gly  
 115 120 125  
 Gln Ile Ile His Gly Asp Arg His Cys Arg Trp Gln Phe Thr Val Glu  
 130 135 140  
 Pro Glu Val Thr Trp Gly Ser Pro Asn Arg Phe Pro Arg Ala Thr Ala  
 35 145 150 155 160  
 Gly Trp Leu Ser Phe Leu Pro Leu Phe Asp Pro Gly Trp Gln Ile Leu  
 165 170 175  
 Leu Ala Gln Gly Arg Ala His Gly Trp Leu Lys Trp Gln Arg Glu Gln  
 180 185 190

Tyr Glu Phe Asp His Ala Leu Val Tyr Ala Glu Lys Asn Trp Gly His  
 .195 200 205  
 Ser Phe Pro Ser Arg Trp Phe Trp Leu Gln Ala Asn Tyr Phe Pro Asp  
 210 215 220  
 5 His Pro Gly Leu Ser Val Thr Ala Ala Gly Gly Glu Arg Ile Val Leu  
 225 230 235 240  
 Gly Arg Pro Glu Glu Val Ala Leu Ile Gly Leu His His Gln Gly Asn  
 245 250 255  
 Phe Tyr Glu Phe Gly Pro Gly His Gly Thr Val Thr Trp Gln Val Ala  
 10 260 265 270  
 Pro Trp Gly Arg Trp Gln Leu Lys Ala Ser Asn Asp Arg Tyr Trp Val  
 275 280 285  
 Lys Leu Ser Gly Lys Thr Asp Lys Lys Gly Ser Leu Val His Thr Pro  
 290 295 300  
 15 Thr Ala Gln Gly Leu Gln Leu Asn Cys Arg Asp Thr Thr Arg Gly Tyr  
 305 310 315 320  
 Leu Tyr Leu Gln Leu Gly Ser Val Gly His Gly Leu Ile Val Gln Gly  
 325 330 335  
 Glu Thr Asp Thr Ala Gly Leu Glu Val Gly Gly Asp Trp Gly Leu Thr  
 20 340 345 350  
 Glu Glu Asn Leu Ser Lys Lys Thr Val Pro Phe  
 355 360

<210> 40  
 25 <211> 56  
 <212> DNA  
 <213> Artificial Sequence

<400> 40  
 30 cgcgatttaa atggcgcgcc ctgcaggcgg ccgcctgcag ggcgcgccat ttaaat 56

<210> 41  
 <211> 32  
 <212> DNA  
 35 <213> Artificial Sequence

<400> 41  
 tcgaggatcc gcggccgcaa gettctctgca gg 32

<210> 42  
<211> 32  
<212> DNA  
<213> Artificial Sequence  
5  
<400> 42  
tcgacctgca ggaagcttgc ggccgcggat cc 32  
  
<210> 43  
10 <211> 32  
<212> DNA  
<213> Artificial Sequence  
  
<400> 43  
15 tcgacctgca ggaagcttgc ggccgcggat cc 32  
  
<210> 44  
<211> 32  
<212> DNA  
20 <213> Artificial Sequence  
  
<400> 44  
tcgaggatcc gcggccgcaa gcttcctgca gg 32  
  
<210> 45  
25 <211> 36  
<212> DNA  
<213> Artificial Sequence  
  
<400> 45  
30 tcgaggatcc gcggccgcaa gcttcctgca ggagct 36  
  
<210> 46  
<211> 28  
35 <212> DNA  
<213> Artificial Sequence  
  
<400> 46  
cctgcaggaa gcttgcggcc gcggatcc 28



<210> 47  
<211> 36  
<212> DNA  
5 <213> Artificial Sequence

<400> 47  
tcgacctgca ggaagcttgc ggccgcggat ccagct 36

10 <210> 48  
<211> 28  
<212> DNA  
<213> Artificial Sequence

15 <400> 48  
ggatccgcgg ccgcaagctt cctgcagg 28

<210> 49  
<211> 39  
20 <212> DNA  
<213> Artificial Sequence

<400> 49  
gatcacctgc aggaagcttg cggccgcgga tccaatgca 39

25 <210> 50  
<211> 31  
<212> DNA  
<213> Artificial Sequence

30 <400> 50  
ttggatccgc ggccgcaagc ttcctgcagg t 31

<210> 51  
35 <211> 41  
<212> DNA  
<213> Artificial Sequence

<400> 51

ggatccgcgg ccgcacaatg gagtctctgc tctctagttc t 41

<210> 52  
<211> 38  
5 <212> DNA  
<213> Artificial Sequence

<400> 52  
ggatcctgca ggtcacttca aaaaaggtaa cagcaagt 38  
10

<210> 53  
<211> 45  
<212> DNA  
<213> Artificial Sequence  
15

<400> 53  
ggatccgcgg ccgcacaatg gcgttttttg ggctctcccg tgttt 45  
20

<210> 54  
<211> 40  
<212> DNA  
<213> Artificial Sequence

<400> 54  
25 ggatcctgca ggttattgaa aacttcttcc aagtacaact 40

<210> 55  
<211> 38  
<212> DNA  
30 <213> Artificial Sequence

<400> 55  
ggatccgcgg ccgcacaatg tggcgaagat ctgttggtt 38  
35

<210> 56  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<400> 56  
ggatcctgca ggcatggag agtagaagga aggagct 37

<210> 57  
5 <211> 50  
<212> DNA  
<213> Artificial Sequence

<400> 57  
10 ggatccgcgg ccgcacaatg gtacttgccg aggttccaaa gcttgctct 50

<210> 58  
<211> 38  
<212> DNA  
15 <213> Artificial Sequence

<400> 58  
ggatcctgca ggtcacttgt ttctggtgat gactctat 38

<210> 59  
20 <211> 38  
<212> DNA  
<213> Artificial Sequence

<400> 59  
25 ggatccgcgg ccgcacaatg acttcgatto tcaacact 38

<210> 60  
<211> 36  
30 <212> DNA  
<213> Artificial Sequence

<400> 60  
35 ggatcctgca ggtcagtgtt gcgatgctaa tgccgt 36

<210> 61  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<400> 61  
taatgtgtac attgtcggcc tc 22

5 <210> 62  
<211> 60  
<212> DNA  
<213> Artificial Sequence

10 <400> 62  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt ccacaattcc ccgcaccgtc 60  
  
<210> 63  
<211> 22  
15 <212> DNA  
<213> Artificial Sequence

<400> 63  
aggctaataa gcacaaatgg ga 22

20 <210> 64  
<211> 63  
<212> DNA  
<213> Artificial Sequence

25 <400> 64  
ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggaattgg tttaggttat 60  
ccc 63

30 <210> 65  
<211> 26  
<212> DNA  
<213> Artificial Sequence

35 <400> 65  
ggatccatgg ttgcccaaac cccatc 26  
  
<210> 66  
<211> 61

<212> DNA  
<213> Artificial Sequence

<400> 66  
5 gcaatgtaac atcagagatt ttgagacaca acgtggcttt gggtaagcaa caatgaccgg 60  
c 61

<210> 67  
<211> 25  
10 <212> DNA  
<213> Artificial Sequence

<400> 67  
gaattctcaa agccagccca gtaac 25

15  
<210> 68  
<211> 63  
<212> DNA  
<213> Artificial Sequence

20  
<400> 68  
ggatgagtc agcaacacct tcttcacgag gcagacctca gcgggtgcga aaagggtttt 60  
ccc 63

25 <210> 69  
<211> 23  
<212> DNA  
<213> Artificial Sequence

30 <400> 69  
ccagtggttt aggtgtgtg gtc 23

<210> 70  
<211> 21  
35 <212> DNA  
<213> Artificial Sequence

<400> 70  
ctgagttgga tgtattggat c 21

<210> 71  
<211> 28  
<212> DNA  
5 <213> Artificial Sequence

<400> 71  
ggatccatgg ttacttcgac aaaaatcc 28

10 <210> 72  
<211> 60  
<212> DNA  
<213> Artificial Sequence

15 <400> 72  
gcaatgtaac atcagagatt ttgagacaca acgtggcttt gctaggcaac cgcttagtac 60

<210> 73  
<211> 28  
20 <212> DNA  
<213> Artificial Sequence

<400> 73  
gaattcttaa cccaacagta aagttccc 28

25 <210> 74  
<211> 63  
<212> DNA  
<213> Artificial Sequence

30 <400> 74  
ggtatgagtc agcaacacct tcttcacgag gcagacctca gcgccggcat tgtctttttac 60  
atg 63

35 <210> 75  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<400> 75  
ggaacccttg cagccgcttc 20

<210> 76  
5 <211> 22  
<212> DNA  
<213> Artificial Sequence

<400> 76  
10 gtatgcccaa ctggtgcaga gg 22

<210> 77  
<211> 28  
<212> DNA  
15 <213> Artificial Sequence

<400> 77  
ggatccatgt ctgacacaca aaataccg 28

<210> 78  
20 <211> 62  
<212> DNA  
<213> Artificial Sequence

<400> 78  
25 gcaatgtaac atcagagatt ttgagacaca acgtggcttt cgccaatacc agccaccaac 60  
ag 62

<210> 79  
30 <211> 27  
<212> DNA  
<213> Artificial Sequence

<400> 79  
35 gaattctcaa atccccgcat ggcctag 27

<210> 80  
<211> 65  
<212> DNA

<213> Artificial Sequence

<400> 80

5 ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggcctacg gcttggacgt 60  
gtggg 65

<210> 81

<211> 21

<212> DNA

10 <213> Artificial Sequence

<400> 81

cacttggatt ccctgatct g 21

15 <210> 82

<211> 21

<212> DNA

<213> Artificial Sequence

20 <400> 82

gcaatacccg cttggaaaac g 21

<210> 83

<211> 29

25 <212> DNA

<213> Artificial Sequence

<400> 83

30 ggatccatga ccgaatcttc gccctagc 29

<210> 84

<211> 61

<212> DNA

<213> Artificial Sequence

35

<400> 84

gcaatgtaac atcagagatt ttgagacaca acgtggettt caatcctagg tagccgaggc 60  
g 61



<210> 85  
<211> 27  
<212> DNA  
<213> Artificial Sequence  
5  
<400> 85  
gaattcttag cccaggccag cccagcc 27  
  
<210> 86  
10 <211> 66  
<212> DNA  
<213> Artificial Sequence  
  
<400> 86  
15 ggtatgagtc agcaacacct tcttcacgag gcagacctca gcggggaatt gatttgttta 60  
attacc 66  
  
<210> 87  
<211> 21  
20 <212> DNA  
<213> Artificial Sequence  
  
<400> 87  
gcgatcgcca ttatcgcttg g 21  
25  
  
<210> 88  
<211> 24  
<212> DNA  
<213> Artificial Sequence  
30  
<400> 88  
gcagactggc aattatcagt aacg 24  
  
<210> 89  
35 <211> 25  
<212> DNA  
<213> Artificial Sequence  
  
<400> 89

ccatggattc gagtaaagtt gtcgc

25

&lt;210&gt; 90

&lt;211&gt; 0

5 &lt;213&gt; Artificial Sequence

&lt;400&gt; 90

gaattcactt caaaaaaggt aacag

10 &lt;210&gt; 91

&lt;211&gt; 4550

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis sp

15 &lt;400&gt; 91

at	tttacacc	aatttgatca	cttaactaaa	ttaattaaat	tagatgatta	tcccaccata	60
t	tttttgagca	ttaaaccata	aaaccatagt	tataagtaac	tgttttaate	gaatatgact	120
c	gattaagat	taggaaaaat	ttataaccgg	taattaagaa	aacattaacc	gtagtaaccg	180
t	aaatgccga	ttcctccctt	gtctaaaaga	cagaaaacat	atattttatt	ttgccccata	240
20	tgtttcactc	tatttaattt	caggcacaat	acttttggtt	ggtaacaaaa	ctaaaaagga	300
ca	acacgtga	tacttttcct	cgtccgtcag	tcagattttt	tttaactag	aaacaagtgg	360
caa	atctaca	ccacattttt	tgtttaatct	attaacttgt	aagttttaaa	ttcctaaaaa	420
agt	ctaacta	attcttctaa	tataagtaca	ttccctaaat	ttcccaaaaa	gtcaaattaa	480
ta	attttcaa	aatctaact	aaatatctaa	taattcaaaa	tcattaaaaa	gacacgcaac	540
25	aatgacacca	attaatcatc	ctcgaccac	acaattctac	agttctcatg	ctaaaccata	600
t	tttttgctc	tctgttcctt	caaaatcatt	tctttctctt	ctttgattcc	caaagatcac	660
tt	ctttgtct	ttgatttttg	at	tttttttc	tctctggcgt	gaaggaagaa	720
at	ggagtcctc	tgctctctag	ttcttctctt	gtttccgctg	gtaaactctg	tccttttctg	780
gt	ttcaggtt	ttatttggtg	tttaggtttc	gtttttgtga	ttcagaacca	tacaaaaagt	840
30	ttgaactttt	ctgaatataa	aataaggaaa	aagtttcgat	ttttataatg	aattgtttac	900
tag	atcgaag	taggtgacaa	aggttattgt	gtggagaagc	ataatttctg	ggcttgactt	960
tga	atttttgt	ttctcatgca	tgcaacttat	caatcagctg	gtgggttttg	ttggaagaag	1020
caga	atctaa	agctccactc	tttatcaggt	tcgttagggg	tttatgggtt	tttgaaatta	1080
aata	ctcaat	catcttagtc	tcattattct	attgggtgaa	tcacattttc	taatttgga	1140
35	tttatgagac	aatgtatggt	ggacttaggt	gaagttcttc	tctttgggtta	tagttgaagt	1200
gt	tactgatg	ttgttttagct	ctttacacca	atatatacac	ccaattttgc	agaaatccga	1260
gt	tctcggtt	gtgattcgag	taaagttgtc	gcaaaaccga	agtttaggaa	caatcttggt	1320
agg	cctgatg	gtcaaggatc	ttcattgttg	ttgtatccaa	aacataagtc	gagatttcgg	1380
gt	taatgcc	ctgcgggtca	gcctgaggct	ttcactcga	atagcaaaca	gaagtctttt	1440

	agagactcgt tagatgcgtt ttacagggtt tctaggcctc atacagttat tggcacagtt	1500
	aagtttctct ttaaaaatgt aactctttta aaacgcaatc tttcagggtt ttcaaggaga	1560
	taacattagc tctgtgattg gatttgcagg tgcttagcat tttatctgta tctttcttag	1620
	cagtagagaa ggtttctgat atatctcctt tacttttcac tggcatcttg gaggtaatga	1680
5	atatataaca cataatgacc gatgaagaag atacattttt ttcgtctctc tgtttaaaca	1740
	attgggtttt gttttcaggc tgttgttgca gctctcatga tgaacattta catagtgtgg	1800
	ctaaatcagt tgtctgatgt tgaaatagat aaggtaacat gcaaattttc ttcatatgag	1860
	ttcgagagac tgatgagatt aatagcagct agtgcctaga tcatctctat gtgggttttt	1920
	gcagggttaac aagccctatc ttccattggc atcaggagaa tattctgtta acaccggcat	1980
10	tgcaatagta gcttccttct ccatcatggg atggtgccat tttcacaaaa tttcaacttt	2040
	tagaattcta taagttagtg aaatagtttg ttataaatcg ttatagagtt tctggcttgg	2100
	gtggattgtt ggttcatggc cattgttctg ggtctttttt gtgagtttca tgctcggtac	2160
	tgcatactct atcaatgtaa gtaagtttct caatactaga atttggctca aatcaaaatc	2220
	tgcagtttct agtttttagg taatgagggt ttaataactt acttctacta caaacagttg	2280
15	ccacttttac ggtggaag atttgcattg gttgcagcaa tgtgtatcct cgctgtccga	2340
	gctattattg ttcaaatcgc cttttatcta catattcagg tactaaacca ttttccttat	2400
	gttttgtagt tgttttcatc aaaatcactt ttatattact aaagctgtga aactttgttg	2460
	cagacacatg tgtttgaag accaatcttg ttcaactagg ctcttatttt cgcactgog	2520
	tttatgagct ttttctctgt cgttattgca ttgtttaagg taaacaaaga tggaaaaaga	2580
20	ttaaatctat gtatacttaa agtaaagcat tctactgtta ttgatgagaa gttttctttt	2640
	ttggttggat gcaggatata cctgatatcg aaggggataa gatattcggg atccgatcat	2700
	tctctgtaac tctgggtcag aaacgggtac gatattctaaa ctaaagaaat tgttttgact	2760
	caagtgttgg attaagatta cagaagaaag aaaactgttt ttgtttcttg caaaattcag	2820
	gtgttttggg catgtgttac actacttcaa atggcttacg ctgttgcaat tctagtgtga	2880
25	gccacatctc cattcatatg gagcaaagtc atctcggtta caatctttct ttacctatcg	2940
	aaaactcgct aattcatcgt ttgagtggta ctggtttcat tttgttccgt tctgttgatt	3000
	ttttttcagg ttgtgggtca tgttatactc gcaacaactt tgtgggctcg agctaagtcc	3060
	gttgatctga gtagcaaaac cgaaataact tcatgttata tgttcatatg gaaggtaga	3120
	ttcgtttata aatagagtct ttactgcctt tttatgcgct ccaatttggg attaaaatag	3180
30	cctttcagtt tcatcgaatc accattatac tgataaattc tcatttctgc atcagctctt	3240
	ttatgcagag tacttgctgt tacctttttt gaagtgactg acattagaag agaagaagat	3300
	ggagataaaa gaataagtca tcactatgct tctgttttta ttacaagttc atgaaattag	3360
	gtagtgaact agtgaattag agttttatc tgaaacatgg cagactgcaa aaatatgtca	3420
	aagatatgaa tttctgttgg gtaaagaagt ctctgcttgg gcaaaatctt aaggttcggg	3480
35	gtgttgatat aatgctaagc gaagaaatcg attctatgta gaaatttccg aaactatgtg	3540
	taaacatgct agaacatctc cattctatat cttcttctgc aagaaagctc tgtttttatc	3600
	acctaaactc tttatctctg tgtagttaag atatgtatat gtacgtgact acattttttt	3660
	gttgatgtaa tttgcagaac gtatggattt ttgttagaaa gcatgagttc gaaagtatat	3720
	gtttatatat atggataatt cagacctaac gtcgaagctc acaagcataa attcactact	3780

	atagtttgc	ctgtaataga	tagttccatt	gatgtcttga	aactgtacgt	aactgcctgg	3840
	gcgttttgtg	gttgatactg	actactgagt	gttctttgtg	agtgttgtaa	gtatacaaga	3900
	agaagaatat	aggctcacgg	gaacgactgt	ggtggaagat	gaaatggaga	tcatcacgta	3960
	gcggttttgc	caaagaccga	gtcacgacg	agtcctatgaa	gtctttacag	ctgctgatta	4020
5	tgattgacca	ttgcttagag	acgcattgga	atcttactag	ggacttgcc	gggagtttct	4080
	tcaagtacgt	gtcagatcat	acgatgtagg	agatttcacg	gctttgatgt	gtttgtttgg	4140
	agtcacaatg	cttaatgggc	ttattggccc	aataatagct	agctcttttg	ctttagccgt	4200
	ttcgtttgtc	ccctgggtgt	gagtattatt	agggtaggt	gtgaccaaag	tcaccagacc	4260
	tagagtgaat	ctagtagagt	cctagacat	ggtccatggc	ttttatttgt	aatttgaaaa	4320
10	atgaacaatt	ctttttgtaa	ggaaaacttt	tatatagtag	acgtttacta	tatagaaact	4380
	agttgaacta	acttcgtgca	attgcataat	aatgggtgtga	aatagagggt	gcaaaaactca	4440
	ataaacattt	cgacgtacca	agagttcgaa	acaataagca	aaatagattt	ttttgcttca	4500
	gactaatttg	tacaatgaat	ggttaataaa	ccattgaagc	ttttattaat		4550
15	<210> 92						
	<211> 4450						
	<212> DNA						
	<213> Arabidopsis sp						
20	<400> 92						
	tttaggttac	aaaatcaatg	atattgcgta	tgtcaactat	aaaagccaaa	agtaaagcct	60
	cttgtttgac	cagaaggta	tgatcattgt	atacatag	ccaaactacc	tcctggaaga	120
	aaagacatgg	atcccaaaca	acaacaatag	cttcttttac	aagaaccagt	agtaactagt	180
	cactaatcta	aaagagttaa	gtttcagctt	ttctggcaat	ggctccttga	tcatttcaat	240
25	cctgaaggag	accactttg	tagcaagacc	atgtcctctg	tttcaactac	agtgtgtctc	300
	aaaagtctac	ttcaattctt	catatatagg	ttcctcacac	tacagcttca	tcctcattcg	360
	ttgacagaga	gagagtcttt	attgaaaact	tcttccaagt	acaactccac	taaatataat	420
	agcaccaaac	cacttgcttg	acacaaatct	gtacagatat	aaaaacacta	ttaggttttc	480
	caaggcaaat	cacataattg	gattgtgaaa	gagtacaaaa	gataaaccca	aattttcata	540
30	ctttctactg	cagtcagcac	cagatgataa	gtcagctgtc	cctatttgcc	atcctaactg	600
	tcctgatgca	gogggcagtg	atgcgtaata	ttgccaccct	taatcattag	agcgagaaac	660
	aaaaagaatc	aaaagacagt	aaatggaatt	aggaatcaca	aatgagtcct	tgtaaagttt	720
	attgagtacc	gagatctgca	ctgaatccag	aaagtgaag	aaaacctatg	gatgctgtgc	780
	caaatccagt	taaccaaagc	tttgatttat	caccgaatct	aagggtgtgt	gacttaacac	840
35	caacttttac	atcatcttct	ttgtctgga	gacacaatat	attagacatt	agtcacatgga	900
	aaaaaatga	tttaacctag	aatatctcaa	aattacttgc	ataaaaactg	aacttgagct	960
	gaaattttgg	gttcgtagct	tgtggcatat	actatttcat	tttcaatggg	ccacaaagggt	1020
	aactttcttt	tctcacttct	gttgcaaagc	ggaagacttt	tatggggcta	actcttcact	1080
	taaagtatag	aaatcagatg	gaaaagggtg	gagatcaggg	taattttctt	ctttatgatt	1140

	gacaaaagtc	gaacatcgaa	atggatgcat	ttgcatgaga	catgaaacaa	aagctgaaaa	1200
	agaaatctgt	ggtggtgaag	ctagaaaaag	aaaacaaagc	aagcaatatg	cacacattga	1260
	gattaactac	tttgctactg	gtcataatca	aatagatttt	gaagctaaaa	aataaaaaag	1320
	gaatatacct	gatgtgcata	aatagtatca	taaacaaggg	tccagcagac	tccggagaga	1380
5	tagagaggga	gtacaataga	tggtgctatg	cttcctttta	ctgcagtcca	tcctaacaat	1440
	gctccccagt	ttatgggtcaa	acctaaaaag	gcttgaggct	gcaattataa	aaacgaatca	1500
	atcataagaa	aatcagaaaa	tatataatgt	ctaactttga	gaagccagaa	tagatttaaa	1560
	ttacccaaaa	tgtaaacctc	ttcataagt	ggtaggaaaa	gacaagtaac	aaagatgaag	1620
	cccctaaaac	acggctgcag	aatatacata	ctgaaatgag	ctcaagtaga	aaagaatttg	1680
10	atcacaaaa	taaagacaag	acctgagaac	atatcttcag	aatttgggcc	aactacataa	1740
	gggtgaacca	tatgtgtatg	tgaattttta	aacaacact	tgcaaatagc	cgactttagg	1800
	gcaagtaaaa	aatccaaaca	aacctgtaat	tgtaaagttg	gagaagaatc	cctaagccta	1860
	aaagcaactg	cagcccgaga	aatccaatcc	cttgaaatgg	tgtcaaaaga	ccactggcga	1920
	taggtcttag	ttttgtacga	tcaacctgga	tataaaagaa	atttgtaaga	caacataatc	1980
15	taaaacaaaa	caaccataca	aaatcttgag	ctttacatac	aagcaacca	tctttgttta	2040
	tggaagaatg	aatccagtta	catgaatgct	gtgtatctac	cctaactact	aaacacatat	2100
	ttcaatcgaa	aaacatattc	caccttcacc	atatctaaca	cctgaagtct	ttcacttttt	2160
	gaacgaagtc	atcagaacat	gcagataagc	tattacccaa	aacagagata	tgactggaaa	2220
	tgttgtcgta	aattgatcca	acatagaaaa	atcaagacca	gttcagatg	tcaaagcaat	2280
20	aacactttcc	caccatgggt	acagaaacca	tagttacaca	aaacatgttt	cctaaaccaa	2340
	catactaaag	ggatatataa	atttgacatc	actttatcac	cataccataa	gatagcttaa	2400
	aaacaaactg	acctttgtat	ctatgtcctg	atcaagcaga	tcatttatag	tacaaccagc	2460
	acctctaaga	agtaatgctc	cgcaaccaa	taaagccata	tattttaa	ttggaaggct	2520
	tccaggatca	gcagccaacg	caatcgacct	atacaacaat	gatggagatt	cagagtatcg	2580
25	atctattttac	atagctctgg	aactagatcc	atgacgaaac	atggaacatc	gttataatat	2640
	ctaaagactt	ccaaacagat	tcctgagtaa	gaaacccagt	ggaactatag	tactgtaaca	2700
	tatataaaat	caaagaaaac	tcagggttat	agcattatcc	aatcctgatt	tctgccaatc	2760
	cttaaccact	ctcccatgct	atcaaaaacc	tcagctcaag	atcatactac	ctaattgcct	2820
	atgagctctt	gggaagatca	ttatggattt	gataactgaa	aaaagtaaca	gagaaatagc	2880
30	agactgcaag	aactactcca	aacttctcca	ctgatatgta	tgtagtctaa	caataataaa	2940
	cagacataaa	ttctttttatc	aagcttcaag	agcaagttag	tcagaaaaca	tcacagccaa	3000
	accaaccagg	aaaacacata	actttatcac	ataaaactaa	atttaagtga	atctgactta	3060
	acataaacca	tcctttggga	cgaaaggaaa	ctatataaac	atgcagtctt	tctttccctc	3120
	agctattctt	tcggatggat	tataatgaat	ctcaaaagtg	aatgtcttg	attctcagct	3180
35	acattactca	aaggcgaaga	taaacttacc	acatacaagg	ccacgcaagc	aaccaagttc	3240
	caatgggttt	atccaatcga	gcaagcttag	cataacctct	aacttcttct	ggtaaataca	3300
	aatctatcca	agaagcttcc	ttaacaacaa	caccatcact	cttctcctta	tcacttttct	3360
	tcggctttcc	ctccaaaacc	gaagaagacg	acgacattcc	acaaattaat	ctgtaattcc	3420
	aaccaacacc	aaaaaacttc	tcctgatgca	attctcttcc	tttactccat	acttggtaat	3480

	tatcattcca tgaaggataa cacttagtga aaggatttgt gtaatgggta gtcacaggat	3540
	tggacaagga tttatgttgt gattgcaaaa gagcagagga agaagatgga gttacggaga	3600
	cggaagattt caacaaccgt cttgaaacac gggagagccc aaaaaacgcc atctttgaga	3660
	gaaattgttg cctggaagaa acaaagactt gagatttcaa acgtaagtga attcttacga	3720
5	acgaaagcta acttctcaag agaatcagat tagtgattcc tcaaaaacaa acaaaactat	3780
	ctaatttcag tttcgagtga tgaagcctta agaactctaga acctccatgg cgtttctaat	3840
	ctctcagaga taatcgaatt ccttaaacaa tcaaagctta gaaagagaag aacaacaaca	3900
	acaacaaaaa aaatcagatt aacaaccgac cagagagcaa cgacgacgcc ggcgagaaag	3960
	agcacgtcgt ctgggagcaa gacttcttct ccagtaacct ggatggatcg ttaatgggcc	4020
10	tgtagattat tatatttggg ccgaaacaat tgggtcagca aaaacttggg ggataatgaa	4080
	gaaacacgta cagtatgcat ttaggctcca aattaattgg ccatataatt cgaatcagat	4140
	aaactaatca acccctacct tacttatttc tcaactgtttt tatttctacc ttagtagttg	4200
	aagaaacact tttatttatc ttttcgggac ccaatttga taggatcggg ccattactca	4260
	tgagcgtcag acacatatta gccttatcag attagtgggg taagggtttt ttaattcggg	4320
15	aagaagcaac aatcaatgct ggagaaatta aagaatctgc atgggcgtgg cgtgatgata	4380
	tgtgcatatg gagtcagttg ccgatcatat ataactattt ataaactaca tataaagact	4440
	actaatagat	4450
	<210> 93	
20	<211> 2850	
	<212> DNA	
	<213> Arabidopsis sp	
	<400> 93	
25	aattaaaaatt tgagcgggtct aaaccattag accgtttaga gatccctcca acccaaaata	60
	gtcgattttc acgtcttgaa catatattgg gccttaactct gtgtgggttag taaagacttt	120
	tatttggtcaa agaaaaacaa ccatggccca acatgttgat acttttattt aattatacaa	180
	gtacccttga attctctgaa atatatttga ttgaccaga tattaatttt aattatcatt	240
	tctgtaaaa gtgaaggagt caccgtgact cgtcgtaatc tgaaccaat ctgttcatat	300
30	gatgaagaag tttctctcgt tctcctccaa cgcgtagaaa attctgacgg cttaacgatg	360
	tggcgaagat ctgttggtta tcgtttctct tcaagaactct ctgtttcttc ttcgttacca	420
	aaccctagac tgattccttg gtcccgcgaa ttatgtgccg ttaatagctt ctccagcct	480
	ccggtctcga cggaatcaac tgctaagtta gggatcactg gtgttagatc tgatgccaat	540
	cgagtttttg ccaactgtac tgcgcgcgt acagctacag ctaccaccgg tgagatttcg	600
35	tctagagttg cggctttggc tggattaggg catcaactacg ctcgttgtta ttgggagctt	660
	tctaaagcta aacttaggta tgtgtttact tttcttttct catgaaaaat ctgaaaattt	720
	ccaattgttg gattcttaaa ttctcatttg ttttatgggt gtagtatgct tgtggttgca	780
	acttctggaa ctgggtatat tctgggtacg ggaaatgctg caattagctt cccggggctt	840
	tgttacacat gtgcaggaac catgatgatt gctgcactg ctaattcctt gaatcaggtc	900

	attgaaatgt	tgagaagttc	ataaatcttcg	aatccttggt	gtgtttatgt	agttgatctt	960
	gcttgcttat	gtttatgtag	ttgaaaagtt	taaaaatttc	taatccttgg	tagttgatct	1020
	cgcttgcttg	ttttttcatt	ttctagattt	ttgagataag	caatgattct	aagatgaaaa	1080
	gaacgatgct	aaggccattg	ccttcaggac	gtattagtgt	tccacacgct	gttgcatggg	1140
5	ctactattgc	tggtgcttct	ggtgcttggt	tggtggccag	caaggtgaat	gtttgttttt	1200
	ttatatgtga	tttctttgtt	ttatgaatgg	gtgattgaga	gattatggat	ctaaactttt	1260
	gcttccacga	caagggttatt	gcagactaat	atgttggtcg	ctggacttgc	atctgccaat	1320
	cttgtacttt	atgcgtttgt	ttatactccg	ttgaagcaac	ttcacccctat	caatacatgg	1380
	gttggtgctg	ttgttggtgc	tatcccaccc	ttgcttggtt	aaatttttgt	tccttttctt	1440
10	ctttatttta	gcagattctg	ttttgttgga	tactgctttt	aattcaaaat	gtagtcatgg	1500
	ttcaccaatt	ctatgcttat	ctatttttgt	tggtgtcagg	tggtggcag	cgtctggtca	1560
	gatttcatac	aattogatga	ttcttccage	tgctctttac	ttttggcaga	tacctcattt	1620
	tatggccctt	gcacatctct	gccgcaatga	ttatgcagct	ggagggtaag	accatatggt	1680
	gtcatatgag	attagaatgt	ctccttccat	gtagtgttga	tcttgaaacta	gttcaatttc	1740
15	gtggaatgat	cagagtgtcc	tagatagtgt	cacagcagtc	gacatttttag	tggttagata	1800
	atgagttcct	tccgttagag	ataaacattc	gcgaacattg	tttccagctt	cgcgcacca	1860
	acttctgatt	ttgtttcttg	gtaccttggt	ttcagttaca	agatgttgtc	actctttgat	1920
	cgcgcagggg	agagaatagc	agcagtggct	ctaaggaaact	gctttttacat	gatccctctc	1980
	ggtttcatcg	cctatgactg	tgagtcttgt	agattcatct	tttttttgta	gtttattgac	2040
20	tgcattgctg	tatctgattt	ttgctgttcc	ttccaatttt	tgtgacaggg	gggttaacct	2100
	caagttgggt	ttgcctcgaa	tcaacacttc	tcacactagc	aatcgctgca	acagcatttt	2160
	cattctaccg	agaccggacc	atgcataaag	caaggaaaat	gttccatgcc	agtcttctct	2220
	tccttctctg	tttcatgtct	ggtcttcttc	tacaccgtgt	ctctaattgat	aatcagcaac	2280
	aactcgtaga	agaagccgga	ttaacaaatt	ctgtatctgg	tgaagtcaaa	actcagaggc	2340
25	gaaagaaacg	tgtggctcaa	cctccggtgg	cttatgcctc	tgctgcaccg	tttcccttcc	2400
	tcccagctcc	ttccttctac	tctccatgat	aacctttaag	caagctattg	aatttttgga	2460
	aacagaaatt	aaaaaaaaa	tctgaaaagt	tcttaagttt	aatcttttgt	taataatgaa	2520
	gtggagaacg	catacaagtt	tatgtatttt	ttctcatctc	cacataattg	tattttttct	2580
	ctaagtatgt	ttcaaattgat	acaaaataca	tactttatca	attatctgat	caaattgatg	2640
30	aattttttgag	ctttgacgtg	ttaggtctat	ctaataaacg	tagtaacgaa	tttggttttg	2700
	gaaatgaaat	ccgataaccg	atgatggtgt	agagttaaac	gattaaaccg	ggttgggttaa	2760
	aggtctcgag	tctcgacggc	tgcggaatc	ggaaaatcac	gattgaggac	tttgagctgc	2820
	cacgaagatg	gcgatgaggt	tgaaatcaat				2850
35	<210> 94						
	<211> 3660						
	<212> DNA						
	<213> Arabidopsis sp						

<400> 94		
	tattttgtatt tttattgtta aattttatga tttcaccggt tatatatcat cccatattaa	60
	tatttagattt attttttggg ctttatttgg gttttcgatt taaactgggc ccattctgct	120
	tcaatgaaac cctaattgggt tttgtttggg ctttggattt aaaccgggccc cattctgctt	180
5	caatgaaggt cctttgtcca acaaaactaa catccgacac aactagtatt gccaaagagga	240
	tcgtgccaca tggcagttat tgaatcaaag gccgccaaaa ctgtaacgta gacattactt	300
	atctccggta acggacaacc actcgtttcc cgaaacagca actcacagac tcacaccact	360
	ccagtctccg gcttaactac caccagagac gattctctct tccgtcgggt ctatgacttc	420
	gattctcaac actgtctcca ccatccactc ttccagagtt acctccgtcg atcgagtcgg	480
10	agtcctctct ctccggaatt cggattccgt tgagttcact cgccggcggt ctggtttctc	540
	gacgttgatc tacgaatcac ccggtagttt gcattctggt ggatagattg atgaatgttt	600
	tcttcgattt ttttttact gatcttggtt tggatctctc gtagggcgga gatttggtgt	660
	gcgtgcggcg gagactgata ctgataaagg tatgattttt tagttgtttt tattttctct	720
	ctcttcaaaa ttctcttttc aaacactgtg gcgtttgaat ttccgacggc agttaaatct	780
15	cagacacctg acaaggcacc agccggtgggt tcaagcatta accagcttct cggatcaaaa	840
	ggagcatctc aagaaactgt aattttgttc atctctcag aatcttttaa attatcatat	900
	ttgtggataa tgatgtgtta gtttaggaat tttctacta aaggtaatct cttttgagga	960
	caagtcttgt ttttagctta gaaatgatgt gaaaatgttg tttgttagct aaaaagagtt	1020
	tggtgttata ttctgtatc agaataaatg gaagattcgt cttcagctta caaaaccagt	1080
20	cacttggcct ccaactggtt ggggagtcgt ctgtggtgct gctgcttcag gtaatcatac	1140
	gaacctcttt tggatcatgc aatactgtac agaaagtttt ttcatcttcc ttccaattgt	1200
	ttctctctgc agggaaactt cattggaccc cagaggatgt tgctaagtcg attctttgca	1260
	tgatgatgtc tggctcctgt ctactggct atacacaggt ctggttttac acaacaaaaa	1320
	gctgacttgt tcttattota gtgcatttgc ttggtgctac aataacctag acttgctgat	1380
25	ttccagacaa tcaacgactg gtatgataga gatatcgacg caattaatga gccatatcgt	1440
	ccaattccat ctggagcaat atcagagcca gaggttaactg agacagaaca ttgtgagctt	1500
	ttatctcttt tgtgattctg atttctcctt actccttaaa atgcagggtta ttacacaagt	1560
	ctgggtgcta ttattgggag gtcttggtat tgctggaata ttagatgtgt gggtaagttg	1620
	gcccttctga cattaactag tacagttaaa gggcacatca gatttgctaa aatcttccct	1680
30	tatcaggcag gccataccac tcccactgtc ttctatcttg ctttgggagg atcattgcta	1740
	tcttatatat actctgctcc acctcttaag gtaagtttta ttccctaact ccaactctcta	1800
	gtgataagac actccatcca agttttggag ttttgaatat cgatatctga actgatctca	1860
	ttgcagctaa aacaaaatgg atgggttggg aattttgcac ttggagcaag ctatattagt	1920
	ttgccatggg aagatatctc gtgtatcaat aatatatggc gttgttctca tctcattgat	1980
35	ttgtttcttg ctcaactgac tgataggtgg gctggccaag cattgttttg cactcttacg	2040
	ccagatgttg ttgttctaac actcttgtag agcatagctg gggtagctct ttggcaaacc	2100
	ttttatgttg cttttttcgt tatctgttgt aatatgctct tgcttcatgt tgtaccttgg	2160
	tgataatgca gtttaggaata gccattgtta acgacttcaa aagtgttgaa ggagatagag	2220
	cattaggact tcagtctctc ccagtagctt ttggcaccga aactgcaaaa tggatatgcg	2280



	ttggtgctat agacattact cagctttctg ttgcgggtat gtactatcca ctgtttttgt	2340
	gcagctgtgg cttctatttc ttttccttga tcttatcaac tggatattca ccaatggtaa	2400
	agcacaaatt aatgaagctg aatcaacaaa ggcaaacat aaaagtacat tctaataaaa	2460
	tgagctaatt aagaggaggc atctactttt atgtttcatt agtggtgattg atggattttc	2520
5	atttcatgct tctaaaacaa gtattttcaa cagtgtcatg aaataacaga acttatatct	2580
	tcatttgtag ttttactagt ggatgagtta cacaatcatt gttatagaac caaatcaaa	2640
	gtagagatca tcattagtag atgtctatct tgggtgcagg atatctatta gcatctggga	2700
	aaccttatta tgcgttggcg ttggttgctt tgatcattcc tcagattgtg ttccaggtaa	2760
	agacgttaac agtctcacat tataattaat caaattcttg tcaactcgtc gattgctaca	2820
10	ctcgtcttcta taaactgcag tttaaatact ttctcaagga ccctgtcaaa tacgacgtca	2880
	agtaccaggc aagtcaactt agtacacatg tttgtgttct tttgaaatat ctttgagagg	2940
	tctcttaatc agaagttgct tgaacactc atcttgatta caggcaagcg cgcagccatt	3000
	cttgggtgctc ggaatatttg taacggcatt agcatcgcaa cactgaaaaa ggcgattttt	3060
	gatgggggtt tgtcgaaagc agaggtgttg acacatcaaa tgtgggcaag tgatggcacc	3120
15	aactagttta aaagattttg taaaatgtat gtaccgttat tactagaaac aactcctgtt	3180
	gtatcaattt agcaaaacgg ctgagaaatt gtaattgatg ttaccgtatt tgcgctccat	3240
	ttttgcattt cctgctcata tcgaggattg gggtttatgt tagttctgtc acttctctgc	3300
	tttcagaatg tttttgtttt ctgtagtggg ttttaactat tttcatcact ttttgtattg	3360
	attctaaaca tgtatccaca taaaacagc aatatacaaa atgatactt cctcaaactt	3420
20	tttataatct aaatctaaca actagctagt aacccaacta acttcataca attaatgtga	3480
	gaaactacaa agactagact atacatatgt tatttaacaa cttgaaactg tgttattact	3540
	acctgatttt tttctattct acagccattt gatatgctgc aatcttaaca tatcaagtct	3600
	cacgttggtg gacacaacat actatcaca gtaagacacg aagtaaaacc aaccggcaac	3660
25	<210> 95	
	<211> 1236	
	<212> DNA	
	<213> SOY	
30	<400> 95	
	atggattcac tgcctctctg atctttccct aatattaata acgcctcttc tctcaccacc	60
	actggtgcaa atttctccag gactaaatct ttgcgcaaca ttaccatgc aagttcttat	120
35	gtgccaaatg cttcatggca caataggaaa atccaaaaag aatataattt tttgagggtt	180
	cgggtggcaa gtttgaacca tcattacaaa ggcattgagg gagcgtgtac atgtaaaaaa	240
	tgtaatatata aatttggtgt gaaagcgacc tctgaaaaat ctcttgagtc tgaacctcaa	300
40	gcttttgatc caaaaagcat tttggactct gtcaagaatt ccttgatgc tttctacagg	360
	ttttccaggc ctcacacagt tattggcaca gcattaagca taatttctgt gtctctctct	420

	getgttgaga aaatatcaga tataatctcca ttatttttta ctggtgtgtt ggaggctgtg	480
5	gttgctgccc tgtttatgaa tatttatatt gttggttga atcaattgtc tgatgttgaa	540
	atagacaaga taaacaagcc gtatcttcca ttagcatctg gggaatattc ctttgaaact	600
	ggtgtcacta ttgttgcac tttttcaatt ctgagttttt ggcttggtctg ggtttaggt	660
10	tcattggccat tattttgggc ctttttgta agctttgtgc taggaactgc ttattcaatc	720
	aatgtgcctc tgttgagatg gaagaggttt gcagtgttg cagcgatgtg cattctagct	780
15	gttcgggcag taatagttca acttgcattt ttccttcaca tgcagactca tgtgtacaag	840
	agggcacctg tcttttcaag accattgatt ttgctactg cattcatgag cttcttctct	900
	gtagttag cactgtttta ggatatacct gacattgaag gagataaagt atttggcatc	960
20	caatcttttt cagtgcgttt aggtcagaag ccggtgttct ggacttgtgt tacccttctt	1020
	gaaatagctt atggagtcgc ctccttggtg ggagctgcat ctccttgtct ttggagcaaa	1080
25	attttcacgg gtctgggaca cgctgtgtg gtttcaattc tctggtttca tgccaaatct	1140
	gtagatttga aaagcaaagc ttcgataaca tccttctata tgtttatttg gaagctattt	1200
	tatgcagaat acttactcat tccttttggt agatga	1236
30		
	<210> 96	
	<211> 1188	
	<212> DNA	
	<213> SOY	
35		
	<400> 96	
	atggattcga tgcttcttcg atcttttctt aatattaaca acgcttcttc tctcgccacc	60
40	actggttctt atttgcaaaa tgcttcatgg cacaatagga aaatccaaaa agaataaat	120
	tttttgaggt ttcggtggcc aagtttgaac caccattaca aaagcattga aggaggtgt	180
	acatgtaaaa aatgtaatat aaaatttggt gtgaaagcga cctctgaaaa atcttttgag	240
45	tctgaacccc aagcttttga tccaaaaagc attttgact ctgtcaagaa ttccttgat	300
	gctttctaca ggttttccag acctcacaca gttattggca cagcattaag cataatttct	360
50	gtgtccctcc ttgctgttga gaaaatatca gatatatctc cattattttt tactggtgtg	420
	ttggaggctg tggttgtgc cctgtttatg aatatttata ttgttggtt gaatcaattg	480
	tctgatgttg aaatagacaa gataaacaag ccgtatcttc cattagcatc tggggaatat	540

tcctttgaaa ctggtgtcac tattgttgca tctttttcaa ttctgagttt ttggcttggc 600  
 tgggtttag gttcatggcc attatatttg gccctttttg taagctttgt gctaggaact 660  
 5 gcttattcaa tcaatgtgcc tctgttgaga tggaagaggt ttgcagtgtc tgcagcgatg 720  
 tgcattctag ctgttcgggc agtaatagtt caacttgcac ttttccttca catccagact 780  
 catgtataca agaggccacc tgtcttttca agatcattga tttttgctac tgcattcatg 840  
 10 agcttcttct ctgtagttaa agcactgttt aaggatatac ctgacattga aggagataaa 900  
 gtatttggca tccaatcttt ttcagtgcgt ttaggtcaga agccggtatt ctggacttgt 960  
 15 gttatccttc ttgaaatagc ttatggagtc gccctcctgg tgggagctgc atctccttgt 1020  
 ctttggagca aaattgtcac ggtctcggga cagctgttc tggttcaat tctctggttt 1080  
 catgccaaat ctgtagattt gaaaagcaaa gcttcgataa catccttcta tatgtttatt 1140  
 20 tggaagctat ttatgcaga atacttactc attccttttg ttagatga 1188

<210> 97

25 <211> 395

<212> PRT

<213> SOY

<400> 97

30

Met Asp Ser Met Leu Leu Arg Ser Phe Pro Asn Ile Asn Asn Ala Ser  
1 5 10 15

Ser Leu Ala Thr Thr Gly Ser Tyr Leu Pro Asn Ala Ser Trp His Asn  
20 25 30

Arg Lys Ile Gln Lys Glu Tyr Asn Phe Leu Arg Phe Arg Trp Pro Ser  
35 40 45

Leu Asn His His Tyr Lys Ser Ile Glu Gly Gly Cys Thr Cys Lys Lys  
50 55 60

Cys Asn Ile Lys Phe Val Val Lys Ala Thr Ser Glu Lys Ser Phe Glu  
65 70 75 80

45

Ser Glu Pro Gln Ala Phe Asp Pro Lys Ser Ile Leu Asp Ser Val Lys  
85 90 95

Asn Ser Leu Asp Ala Phe Tyr Arg Phe Ser Arg Pro His Thr Val Ile  
100 105 110

50

Gly Thr Ala Leu Ser Ile Ile Ser Val Ser Leu Leu Ala Val Glu Lys  
115 120 125

Ile Ser Asp Ile Ser Pro Leu Phe Phe Thr Gly Val Leu Glu Ala Val  
 130 135 140  
 5 Val Ala Ala Leu Phe Met Asn Ile Tyr Ile Val Gly Leu Asn Gln Leu  
 145 150 155 160  
 Ser Asp Val Glu Ile Asp Lys Ile Asn Lys Pro Tyr Leu Pro Leu Ala  
 165 170 175  
 10 Ser Gly Glu Tyr Ser Phe Glu Thr Gly Val Thr Ile Val Ala Ser Phe  
 180 185 190  
 Ser Ile Leu Ser Phe Trp Leu Gly Trp Val Val Gly Ser Trp Pro Leu  
 195 200 205  
 15 Phe Trp Ala Leu Phe Val Ser Phe Val Leu Gly Thr Ala Tyr Ser Ile  
 210 215 220  
 Asn Val Pro Leu Leu Arg Trp Lys Arg Phe Ala Val Leu Ala Ala Met  
 225 230 235 240  
 Cys Ile Leu Ala Val Arg Ala Val Ile Val Gln Leu Ala Phe Phe Leu  
 245 250 255  
 25 His Ile Gln Thr His Val Tyr Lys Arg Pro Pro Val Phe Ser Arg Ser  
 260 265 270  
 Leu Ile Phe Ala Thr Ala Phe Met Ser Phe Phe Ser Val Val Ile Ala  
 275 280 285  
 Leu Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Lys Val Phe Gly Ile  
 290 295 300  
 35 Gln Ser Phe Ser Val Arg Leu Gly Gln Lys Pro Val Phe Trp Thr Cys  
 305 310 315 320  
 Val Ile Leu Leu Glu Ile Ala Tyr Gly Val Ala Leu Leu Val Gly Ala  
 325 330 335  
 40 Ala Ser Pro Cys Leu Trp Ser Lys Ile Val Thr Gly Leu Gly His Ala  
 340 345 350  
 Val Leu Ala Ser Ile Leu Trp Phe His Ala Lys Ser Val Asp Leu Lys  
 355 360 365  
 Ser Lys Ala Ser Ile Thr Ser Phe Tyr Met Phe Ile Trp Lys Leu Phe  
 370 375 380  
 50 Tyr Ala Glu Tyr Leu Leu Ile Pro Phe Val Arg  
 385 390 395  
 <210> 98  
 <211> 411  
 55 <212> PRT  
 <213> SOY

&lt;400&gt; 98

5 Met Asp Ser Leu Leu Leu Arg Ser Phe Pro Asn Ile Asn Asn Ala Ser  
 1 5 10 15  
 Ser Leu Thr Thr Thr Gly Ala Asn Phe Ser Arg Thr Lys Ser Phe Ala  
 20 25 30  
 10 Asn Ile Tyr His Ala Ser Ser Tyr Val Pro Asn Ala Ser Trp His Asn  
 35 40 45  
 Arg Lys Ile Gln Lys Glu Tyr Asn Phe Leu Arg Phe Arg Trp Pro Ser  
 50 55 60  
 15 Leu Asn His His Tyr Lys Gly Ile Glu Gly Ala Cys Thr Cys Lys Lys  
 65 70 75 80  
 Cys Asn Ile Lys Phe Val Val Lys Ala Thr Ser Glu Lys Ser Leu Glu  
 85 90 95  
 20 Ser Glu Pro Gln Ala Phe Asp Pro Lys Ser Ile Leu Asp Ser Val Lys  
 100 105 110  
 25 Asn Ser Leu Asp Ala Phe Tyr Arg Phe Ser Arg Pro His Thr Val Ile  
 115 120 125  
 Gly Thr Ala Leu Ser Ile Ile Ser Val Ser Leu Leu Ala Val Glu Lys  
 130 135 140  
 30 Ile Ser Asp Ile Ser Pro Leu Phe Phe Thr Gly Val Leu Glu Ala Val  
 145 150 155 160  
 Val Ala Ala Leu Phe Met Asn Ile Tyr Ile Val Gly Leu Asn Gln Leu  
 165 170 175  
 35 Ser Asp Val Glu Ile Asp Lys Ile Asn Lys Pro Tyr Leu Pro Leu Ala  
 180 185 190  
 40 Ser Gly Glu Tyr Ser Phe Glu Thr Gly Val Thr Ile Val Ala Ser Phe  
 195 200 205  
 Ser Ile Leu Ser Phe Trp Leu Gly Trp Val Val Gly Ser Trp Pro Leu  
 210 215 220  
 45 Phe Trp Ala Leu Phe Val Ser Phe Val Leu Gly Thr Ala Tyr Ser Ile  
 225 230 235 240  
 Asn Val Pro Leu Leu Arg Trp Lys Arg Phe Ala Val Leu Ala Ala Met  
 245 250 255  
 50 Cys Ile Leu Ala Val Arg Ala Val Ile Val Gln Leu Ala Phe Phe Leu  
 260 265 270  
 55 His Met Gln Thr His Val Tyr Lys Arg Pro Pro Val Phe Ser Arg Pro  
 275 280 285

Leu Ile Phe Ala Thr Ala Phe Met Ser Phe Phe Ser Val Val Ile Ala  
 290 295 300

5 Leu Phe Lys Asp Ile Pro Asp Ile Glu Gly Asp Lys Val Phe Gly Ile  
 305 310 315 320

Gln Ser Phe Ser Val Arg Leu Gly Gln Lys Pro Val Phe Trp Thr Cys  
 325 330 335

10 Val Thr Leu Leu Glu Ile Ala Tyr Gly Val Ala Leu Leu Val Gly Ala  
 340 345 350

15 Ala Ser Pro Cys Leu Trp Ser Lys Ile Phe Thr Gly Leu Gly His Ala  
 355 360 365

Val Leu Ala Ser Ile Leu Trp Phe His Ala Lys Ser Val Asp Leu Lys  
 370 375 380

20 Ser Lys Ala Ser Ile Thr Ser Phe Tyr Met Phe Ile Trp Lys Leu Phe  
 385 390 395 400

Tyr Ala Glu Tyr Leu Leu Ile Pro Phe Val Arg  
 405 410

25

<210> 99  
 <211> 964  
 <212> DNA  
 <213> RICE

30

<400> 99

gagcagcact gggctcttaca ttccaatgga gctcgccctgt tgcttttcatt acatgcttcg 60  
 tgacttttatt tgcttttggtc attgctataa ccaaagatct cccagatggt gaaggggatc 120  
 35 ggaagtatca aatatcaact ttggcgacaa agctcggtgt cagaaacatt gcatttcttg 180  
 gctctgggtt attgatagca aattatgttg ctgctattgc thtagctttt ctcatgcctc 240  
 40 aggcttttcag gcgcactgta atgggtgcctg tgcattgctgc ccttgccgtt ggtataattt 300  
 tccagacatg ggttctggag caagcaaaat atactaagga tgctatttca cagtactacc 360  
 ggttcatttg gaatctcttc tatgctgaat acatcttctt cccgttgata tagagaccaa 420  
 45 gcaatctgat atgggtctgca tgttgagtgc ggcaaaaact agaagcccat atgaacagtg 480  
 ggagtagggg aacgaacatg ccatccatgg gaagactctg ataactctct ctgccccggg 540  
 50 ctgtaaaggg taagcactgt tgggcatata tatgaaagga aggtgataaa gcaggggatgc 600  
 taaattgcta ctgggatcct caaaggctta tagtggtcac cagtggaatg tgccttaata 660  
 atttggttac ccagcagagc aagtttttgc aggttattag gtaatatctt tgagggaatg 720

aacttagatt tcattgtttt aaggtctggt cacacaacgg gtagtagtgc tggagcggca 780  
 5 aaaaacgacc ttgttttaca ctaccaaggg aggttaactc tagttttcat gtgaccactt 840  
 accttgagag ttgagaccat ggaatcactt gtcgactcct cggcttgat atttctagtg 900  
 tcagcatttg cattctcttc cccacttgta cttgaaaagt tgaagacaac ttttttgttt 960  
 10 gtgt 964

<210> 100  
 <211> 421  
 15 <212> DNA  
 <213> WHEAT

<400> 100  
 20 cgtccgcgga cgcgtgggtg cttattcagt caatctgccg cactttctat ggaagagatc 60  
 tgctgttggt gcagcaactct gcatattagc agtgcgtgcg gtgatagtgc aactggcatt 120  
 ttttctccac attcagacat ttgttttcag aaggccggca gacttttcaa agccattgat 180  
 25 atttgcaact gccttcatga cattcttctc agttgtaata gcattattca aggatatacc 240  
 cgatattgaa ggggaccgca tctttggaat ccaatctttt agtggtagac taggtcaaag 300  
 caggggtttc tggacttgcg ttggcctact tgagggtgcc tacggtgttg cgatactgag 360  
 30 gggggtaact tcttcagtt tgtggagcaa atctataact gttgtgggcc atgcaatcct 420  
 c 421

35  
 <210> 101  
 <211> 705  
 <212> DNA  
 <213> LEEK

40  
 <400> 101  
 gtttcccccc ctgaatttt tttttttttt ttttacttca tttttctgtg aataaattct 60  
 taaaaaagac aaagaaaacc actggatatc ctaaattcaa cataggctat tgtcattcaa 120  
 45 tgataatctt taacacaaca tacaacatga atataattaa ggagaaatga tctgcaattg 180  
 ttgaaagaac tctccgtttt taagatgaca attaaagcgt tgtaattcc agccatttct 240  
 50 gcctccatta tctactcatc ttctcttgcg attcttttcc atgtaggtca taaaccctca 300

tcttacaaaa ggaatgagca agtactcagc atagaagagc ttccacacga acatataaaa 360  
 5 agatgtaata gtggttttgg tcattggtcc atgagatcta gcacgattcc aaagtaacga 420  
 cccaagaatt gcatgaccta tcaactgttaa gcatttgctc cataggcatg aggaagtagc 480  
 tccaacaacc atgacaacag tgtaggccat ctcaaggaga tatatacata tccaaaacac 540  
 10 cctctcctgg ccaaggcgca cgctgaaaga atggatgcca aatattttgt ctccgtctat 600  
 atcagggtata tccttaaata gagcaataac aactgagaag aagctcatga aggagttgc 660  
 15 aaatatcaat ggccttgtga aacttgctgg tcttttgaaa acaaa 705

<210> 102  
 <211> 637  
 <212> DNA  
 20 <213> LEBK

<220>  
 <221> misc\_feature  
 <222> (1)..(637)  
 25 <223> n = g, a, t or c

<400> 102  
 30 nattcggcac gagttttgaa gaagttaagc atggactccc tccttaccaa gccagttgta 60  
 atacctctgc cttctccagt ttgttcaact ccaatcttgc gaggcagttc tgcaccaggg 120  
 cagtattcat gtagaaacta caatccaata agaattcaaa ggtgcctcgt aaattatgaa 180  
 35 catgtgaaac caaggtttac aacatgtagt aggtctcaaa aacttggtca tgtaaaagcc 240  
 acatccgagc attctttaga atctggatcc gaaggataca ctctagaag catatgggaa 300  
 40 gccgtactag ctctactgaa tgttctatac aaattttcac gacctcacac aataatagga 360  
 acagcaatgg gcataatgac agtttctttg cttgttgctg agagcctatc cgatatttct 420  
 cctctgtttt ttgtgggatt attagaggct gtggttgctg cattgtttat gaatgtttac 480  
 45 attgtaggtc tgaatcaatt atttgacata gaaatagaca aggtcaataa acctgatctt 540  
 cctcttgcat ctggagaata ctaccaaga gctggtagct ctattgtcat tgcttcagcc 600  
 50 atcatgagct ttggcattgg atggttagtt ggctctt 637



<210> 103  
 <211> 677  
 <212> DNA  
 5 <213> CANOLA

<400> 103  
 tttttttttt tttttttcaa aaagaccaat ccttttagtat gtacatgaac aaagtgattt 60  
 10 tgtctccaag ctacaaagaa gaagaagaga ggtatacaaa gaaaactaca aatgttcacc 120  
 atgaatgcta gaagaagggg aataacagat actctgcgta gaagagattc catataaacc 180  
 15 ggtaatatcc tgctatagct tcctttgtgt agtttgcttt ttctagcacc catgtctgga 240  
 aaaccaagca tgaagccaag atcatatgtg caggaatcat caagctacct ctaaaaacct 300  
 gaggcattgta gaaagctagt gatatggcag aaatatagtt cactagcaga agtccagaac 360  
 20 cgaggaatgc aatgttcctc actccaagct ttgttgctag tgttgatatt tggaaactgc 420  
 gatctccttc aacatcagga agatcttttg taatagcaat gactagtgc aacagtggtca 480  
 caaaagacgt gatgaaagcc acaggtgcac tccactgaaa cgaaagtcca agagcagctc 540  
 25 tagtagcatg gtacacacca aaattaagaa gaaaacctcg taccgtggca ataataagaa 600  
 acgctgcaac tggaaatctc ttcattctaa atgggtggaac agaatagatg gtccccagat 660  
 30 cggacgcgtg ggtcgac 677

<210> 104  
 <211> 1431  
 35 <212> DNA  
 <213> CORN

<400> 104  
 40 ccacgcgtcc gcccggccaa gggatggacg cgcttcgcct acggccgtcc ctctccccg 60  
 tgcgcccgcg cgcgcccgcc cgcgagatc attttctacc accatgttgt tccatacaac 120  
 gaaatggtga aggacgaatt tgcttttcta gccaaaggac ccaaggctcc accttgcatc 180  
 45 accatcagaa attcttcgaa tggaaatcct cctattgtag gatatcacat cggtcattaa 240  
 atacttctgt taatgcttcg gggcaacagc tgcagtctga acctgaaaca catgattcta 300  
 50 caaccatctg gagggcaata tcatcttctc tagatgcatt ttacagattt tcccggccac 360

atactgtcat aggaacagca ttaagcatag tctcagtttc ccttctagct gtccagagct 420  
 tgtctgatat atcacctttg ttcctcactg gtttgctgga ggcagtggta gctgcccttt 480  
 5 tcatgaatat ctatattgtt ggactgaacc agttattcga cattgagata gacaaggtta 540  
 acaagccaac tcttccattg gcatctgggg aatacaccct tgcaactggg gttgcaatag 600  
 10 tttcggctct tgccgctatg agctttggcc ttggatgggc tgttggatca caacctctgt 660  
 tttgggctct tttcataagc tttgttcttg ggactgcata ttcaatcaat ctgccgtacc 720  
 ttcgatggaa gagatttget gttgttgacg cactgtgcat attagcagtt cgtgcagtga 780  
 15 ttgttcagct ggcctttttt ctccacattc agacttttgt tttcaggaga ccggcagtgt 840  
 tttctaggcc attattattt gcaactggat ttatgacgtt cttctctgtt gtaatagcac 900  
 tattcaagga tatacctgac atogaagggg accgcatatt cgggatccga tccttcagcg 960  
 20 tccggttagg gcaaaagaag gtcttttggg tctgcgttg cttgcttgag atggcctaca 1020  
 gcgttgcat actgatggga gctacctctt cctgtttgtg gagcaaaaca gcaaccatcg 1080  
 25 ctggccattc catacttgcc gcgatctat ggagctgcgc gcgatcggcg gacttgacga 1140  
 gcaaagccgc aataacgtcc ttctacatgt tcatctggaa gctgttctac gcggagtacc 1200  
 30 tgctcatccc tctggtgcgg tgagcgcgag gcgaggtggt ggcagacgga tcggcgctcg 1260  
 cggggcggca aacaactcca cgggagaact tgagtgcggg aagtaaaact ccgtttgaaa 1320  
 gttgaagcgt gcaccaccgg caccgggcag agagagacac ggtggctgga tggatacggg 1380  
 35 tggccccccc aataaattcc cccgtgcatg gtaaaaaaaaa aaaaaaaaaa a 1431

&lt;210&gt; 105

&lt;211&gt; 1870

40 &lt;212&gt; DNA

&lt;213&gt; CORN

&lt;400&gt; 105

gccgcgcagc gcgacgagcg ccaactgctt gctgccgctt gcctgcgtgc gtgtgcgtcc 60  
 45 accactgacc ccgcgcccgc ccgcgcccc tgccctcca ctccacttgc tcaactgctg 120  
 cggcccgtt ccccccgcc caagggatgg acgcgcttcg cctacggccg tccctcctcc 180  
 50 ccgtgcggcc cggcgcggcc cgcgcgcgag atcattttct accaccatgt tgttccatac 240  
 aacgaaatgg tgaaggacga atttgctttt ctagccaaag gaccaaggt cctaccttgc 300  
 atcaccatca gaaattcttc gaatggaaat cctcctattg taggatatca catcggtcat 360

	taaataacttc tgttaatgct tcggggcaac agctgcagtc tgaacctgaa acacatgatt	420
	ctacaaccat ctggagggca atatcatctt ctctagatgc attttacaga ttttcccggc	480
5	cacatactgt cataggaaca gcattaagca tagtctcagt ttcccttcta gctgtccaga	540
	gcttgtctga tatatcacct ttgttctca ctggtttgct ggaggcagtg gtagctgcc	600
10	ttttcatgaa tatctatatt gttggactga accagttatt cgacattgag atagacaagg	660
	ttaacaagcc aactcttcca ttggcatctg gggaatacac ccttgcaact ggggttgcaa	720
	tagtttcggt ctttgccgct atgagctttg gccttggatg ggctgttgga tcacaacctc	780
15	tgttttgggc tcttttcata agctttgttc ttgggactgc atattcaatc aatctgccgt	840
	accttcgatg gaagagattt gctgttgttg cagcactgtg catattagca gttcgtgcag	900
20	tgattgttca gctggccttt tttctccaca ttcagacttt tgttttcagg agaccggcag	960
	tgttttctag gccattatta ttgcaactg gatttatgac gttcttctct gttgtaatag	1020
	cactattcaa ggatatacct gacatcgaag gggaccgcat attcgggatc cgatccttca	1080
25	gcgtccggtt agggcaaaag aaggctcttt ggatctgcgt tggcttgctt gagatggcct	1140
	acagcgttgc gatactgatg ggagctacct cttcctgttt gtggagcaaa acagcaacca	1200
30	tcgctggcca ttccatactt gccgcgatcc tatggagctg cgcgcgatcg gtggacttga	1260
	cgagcaaagc cgcaataacg tcttctaca tgttcatctg gaagctgttc tacgcggagt	1320
	acctgctcat ccctctggtg cggtgagcgc gaggcgaggt ggtggcagac ggatcggcgt	1380
35	cggcggggcg gcaaacaact ccacgggaga acttgagtgc cggaagtaaa ctcccgtttg	1440
	aaagtgaag cgtgcaccac cggcaccggg cagagagaga cacggtggct ggatggatac	1500
40	ggatggcccc cccaataaat tccccgtgc atggtacccc acgctgcttg atgatatccc	1560
	atgtgtccgg gtgaccggac ctgatcgtct ctagagagat tggttgcaca acgtccaaca	1620
	tagcccgtag gtattgctac cactgctagt atgatactcc ttcctagtcc ttgccagcac	1680
45	cagtgaacca aacttggtcg gctgagctca gcgctcagca gctttacgtg catctgcgcc	1740
	ttgacttgtg cagtgggcgt cgctagcatg aatgatgtat ggtgcgtcac ggctgacgg	1800
50	ttcgtcagtc tgggcctgtt tttgtgtccg aggaagatcg tctgtcagag atctggattg	1860
	cctcgtgct	1870
55	<210> 106	
	<211> 642	

&lt;212&gt; DNA

&lt;213&gt; CORN

&lt;400&gt; 106

5 cggccggact cttctgactt ggcaaccgcc ggcagcgcg acgagcgcca cctgcttgct 60  
 gccgcgtgcc tgcgtgcgtg tgcgtccacc actgaccccg cgcgcgcgcg ccgcccctgc 120  
 10 ccctccactc cacttgctca ctgcgcggct cgtcgcggcc cgcttcccc ccggccaagg 180  
 gatggacgcg cttgcctac gccgcgcct cctccccgtg cggcccggcg cggcccgccc 240  
 gcgaggcagt gtagctgcc cttttcatga atatctatat tggtggactg aaccagttat 300  
 15 tcgacattga gatagacaag gttaacaagc caactcttcc attggcatct ggggaataca 360  
 cccttgcaac tggggttgca atagtttcgg tctttgcgc tatgagcttt ggccttggat 420  
 gggctgttgg atcacaacct ctgttttggg ctcttttcat aagctttgtt cttgggactg 480  
 20 catattcaat caatctgccg taccttcgat ggaagagatt tgctgttggt gcagcactgt 540  
 gcatattagc agttcgtgca gtgattgttc agctggcctt tttctccac attcagactt 600  
 25 ttgttttcag gagaccggca gtgttttcta ggccattatt at 642

&lt;210&gt; 107

&lt;211&gt; 362

30 &lt;212&gt; DNA

&lt;213&gt; COTTON

&lt;400&gt; 107

35 cccacgcgtc cgaacattgt ttgcacttgt tattgccata accaaggatc ttccagatgt 60  
 agaaggagat cgcaaatttc aaatatcaac attagcaaca aagcttggag ttagaaatat 120  
 tgcatttctt ggttcggac ttctactggt gaattatgtt gctgctgtgt tggtgcaat 180  
 40 atacatgcct caggctttca gccgtagttt aatgatacct gctcatatct ttttggcggt 240  
 ctgcttgatt tttcagacat ggggtgttga acaagcaaat tacaaaaagg aagcaatctc 300  
 45 ggggttctat cgtttcatat ggaatctctt ctatgcagag tatgcgattt tccccttcgt 360  
 gt 362

&lt;210&gt; 108

50 &lt;211&gt; 575

&lt;212&gt; DNA

&lt;213&gt; TOMATO

&lt;400&gt; 108

5 cagatcaatt ccagttcctg ctgagttttc tccactcaaa accagttcac atgcaatagt 60  
 acgggttttg aaatgtaaag catggaagag accaaaaaag cactattcct cttcaatgaa 120  
 gttgcagcgg cagtatatca cgcaagagca tgttgaggga agtgatctaa gcactattgc 180  
 10 tgctgataaa aaacttaaag ggagattttt ggtgcacgca tcatctgaac accctcttga 240  
 atctcaacct tctaaaagtc cttgggactc agttaatgat gccgtagatg ctttctacag 300  
 15 gttctcgcgg ccccatacca taataggaac agcattgagc ataatttcag tttctctcct 360  
 tgcagttgag aagttctctg atttttctcc attatttttc actgggggtgt tagaggccat 420  
 tgttgctgcc ctattcatga acatttacat agttggttta aaccagttgt ctgacatcga 480  
 20 aatagacaag gtaacaagc catatcttcc attggcatca ggggaatact ctgtacaaac 540  
 tggagtgatt gttgtgtcgt cttttgccat ttgga 575

25

&lt;210&gt; 109

&lt;211&gt; 1663

&lt;212&gt; DNA

&lt;213&gt; ARABIDOPSIS

30

&lt;400&gt; 109

aacaccaaac acacaatttc acattctttt gcattttct tcttcttctt ccattatgga 60  
 gatacggagc ttgattgttt ctatgaacct taatttatct tcctttgagc tctctcgccc 120  
 35 tgtatctcct ctcactcgct cactagttcc gttecgatcg actaaactag ttccccgctc 180  
 catttctagg gggatcccggt cgatctccac ccgaatagt gaaactgaca agatctccgt 240  
 40 taaacctgtt tacgtccga cgtctcccaa tcgcgaactc cggactctc acagtggata 300  
 ccatttcgat ggaacacctc ggaagttctt cgagggatgg tggatccggg tttccatccc 360  
 agagaagagg gagagttttt gttttatgta ttctgtggag aatcctgcat ttcggcagag 420  
 45 tttgtcacca ttggaagtgg ctctatatgg acctagattc actggtgttg gagctcagat 480  
 tcttggcgct aatgataaat atttatgcc aacgaacaa gactctcaca atttctgggg 540  
 50 agatcgacat gagctagttt tggggaatac ttttagtgct gtgccaggcg caaaggctcc 600  
 aaacaaggag gttccaccag aggaatttaa cagaagagtg tccgaagggt tccaagctac 660

tccatttttg catcaaggtc acatttgcca tgatggccgt actgactatg cggaaactgt 720  
 5 gaaatctgct cggtgggagt atagtactcg tcccgtttac ggttggggtg atgttggggc 780  
 caaacagaag tcaactgcag gctggcctgc agcttttctt gtatttgagc ctcatgggca 840  
 gatatgcatg gcaggaggcc tttccacagg gtggatagaa tggggcggtg aaaggtttga 900  
 10 gtttcgggat gcaccttctt attcagagaa gaattggggt ggaggcttcc caagaaaatg 960  
 gttttgggtc cagtgtaatg tctttgaagg ggcaactgga gaagttgctt taaccgcagg 1020  
 tggcgggttg aggcaattgc ctggattgac tgagacctat gaaaatgctg cactggtttg 1080  
 15 tgtacactat gatggaaaaa tgtacgagtt tgttccttgg aatggtgttg ttagatggga 1140  
 aatgtctccc tggggttatt ggtatataac tgcagagAAC gaaaaccatg tgggtggaact 1200  
 20 agaggcaaga acaaatgaag cgggtacacc tctgcgtgct cctaccacag aagttgggct 1260  
 agctacggct tgcagagata gttgttacgg tgaattgaag ttgcagatat gggaacggct 1320  
 atatgatgga agtaaaggca aggtgatatt agagacaaag agtcaatgg cagcagtgga 1380  
 25 gataggagga ggaccgtggt ttgggacatg gaaaggagat acgagcaaca cggccgagct 1440  
 actaaaacag gctcttcagg tccattgga tcttgaaagc gccttaggtt tggtcctttt 1500  
 30 cttcaagcca ccgggtctgt aacattgatg agtgttttgt ttgttgatag agacccatgt 1560  
 gatgaatgaa gccttagtca tgcattgct agcttcaacta ttatgtatgt atgattttag 1620  
 35 ttcgttcggt ccttgtggta aatgatacgg gccagtgtaa agt 1663

&lt;210&gt; 110

&lt;211&gt; 488

&lt;212&gt; PRT

40 &lt;213&gt; ARABIDOPSIS

&lt;400&gt; 110

45 Met Glu Ile Arg Ser Leu Ile Val Ser Met Asn Pro Asn Leu Ser Ser  
 1 5 10 15  
 Phe Glu Leu Ser Arg Pro Val Ser Pro Leu Thr Arg Ser Leu Val Pro  
 20 25 30  
 50 Phe Arg Ser Thr Lys Leu Val Pro Arg Ser Ile Ser Arg Val Ser Ala  
 35 40 45  
 Ser Ile Ser Thr Pro Asn Ser Glu Thr Asp Lys Ile Ser Val Lys Pro

	50		55		60	
	Val Tyr Val Pro Thr Ser	Pro Asn Arg Glu Leu Arg Thr Pro His Ser				
	65	70	75	80		
5	Gly Tyr His Phe Asp Gly Thr Pro Arg Lys Phe Phe Glu Gly Trp Tyr					
		85	90	95		
10	Phe Arg Val Ser Ile Pro Glu Lys Arg Glu Ser Phe Cys Phe Met Tyr					
		100	105	110		
	Ser Val Glu Asn Pro Ala Phe Arg Gln Ser Leu Ser Pro Leu Glu Val					
		115	120	125		
15	Ala Leu Tyr Gly Pro Arg Phe Thr Gly Val Gly Ala Gln Ile Leu Gly					
		130	135	140		
	Ala Asn Asp Lys Tyr Leu Cys Gln Tyr Glu Gln Asp Ser His Asn Phe					
		145	150	155	160	
20	Trp Gly Asp Arg His Glu Leu Val Leu Gly Asn Thr Phe Ser Ala Val					
		165	170	175		
	Pro Gly Ala Lys Ala Pro Asn Lys Glu Val Pro Pro Glu Glu Phe Asn					
		180	185	190		
25	Arg Arg Val Ser Glu Gly Phe Gln Ala Thr Pro Phe Trp His Gln Gly					
		195	200	205		
30	His Ile Cys Asp Asp Gly Arg Thr Asp Tyr Ala Glu Thr Val Lys Ser					
		210	215	220		
	Ala Arg Trp Glu Tyr Ser Thr Arg Pro Val Tyr Gly Trp Gly Asp Val					
		225	230	235	240	
35	Gly Ala Lys Gln Lys Ser Thr Ala Gly Trp Pro Ala Ala Phe Pro Val					
		245	250	255		
	Phe Glu Pro His Trp Gln Ile Cys Met Ala Gly Gly Leu Ser Thr Gly					
		260	265	270		
40	Trp Ile Glu Trp Gly Gly Glu Arg Phe Glu Phe Arg Asp Ala Pro Ser					
		275	280	285		
45	Tyr Ser Glu Lys Asn Trp Gly Gly Gly Phe Pro Arg Lys Trp Phe Trp					
		290	295	300		
	Val Gln Cys Asn Val Phe Glu Gly Ala Thr Gly Glu Val Ala Leu Thr					
		305	310	315	320	
50	Ala Gly Gly Gly Leu Arg Gln Leu Pro Gly Leu Thr Glu Thr Tyr Glu					
		325	330	335		
	Asn Ala Ala Leu Val Cys Val His Tyr Asp Gly Lys Met Tyr Glu Phe					
		340	345	350		
55	Val Pro Trp Asn Gly Val Val Arg Trp Glu Met Ser Pro Trp Gly Tyr					
		355	360	365		

Trp Tyr Ile Thr Ala Glu Asn Glu Asn His Val Val Glu Leu Glu Ala  
370 375 380

5 Arg Thr Asn Glu Ala Gly Thr Pro Leu Arg Ala Pro Thr Thr Glu Val  
385 390 395 400

Gly Leu Ala Thr Ala Cys Arg Asp Ser Cys Tyr Gly Glu Leu Lys Leu  
405 410 415

10 Gln Ile Trp Glu Arg Leu Tyr Asp Gly Ser Lys Gly Lys Val Ile Leu  
420 425 430

Glu Thr Lys Ser Ser Met Ala Ala Val Glu Ile Gly Gly Gly Pro Trp  
15 435 440 445

Phe Gly Thr Trp Lys Gly Asp Thr Ser Asn Thr Pro Glu Leu Leu Lys  
450 455 460

20 Gln Ala Leu Gln Val Pro Leu Asp Leu Glu Ser Ala Leu Gly Leu Val  
465 470 475 480

Pro Phe Phe Lys Pro Pro Gly Leu  
25 485